arXiv:1003.1781v4 [math.SG] 1 Dec 2011

TOPOLOGICAL QUANTUM FIELD THEORY STRUCTURE ON SYMPLECTIC COHOMOLOGY

ALEXANDER F. RITTER

Abstract. We construct the TQFT on twisted symplectic cohomology and twisted wrapped Floer cohomology, and prove that Viterbo restriction preserves the TQFT. This yields new applications in symplectic topology relating to the Arnol'd chord conjecture and to exact contact embeddings. In the Appendix we construct in detail the untwisted TQFT theory, which leads to a new result: if a Liouville domain M admits an exact embedding into an exact convex symplectic manifold X, and the boundary ∂M is displaceable in X, then the symplectic cohomology of M vanishes and the chord conjecture holds for any Lagrangianly fillable Legendrian in ∂M. The TQFT respects the isomorphism between the symplectic cohomology of a cotangent bundle and the homology of the free loop space, so it recovers the TQFT of string topology.

0.1. **Motivation.** Symplectic cohomology has become an important tool in symplectic topology. For example, it can be used to prove the existence of Hamiltonian orbits and Reeb chords, and it gives rise to obstructions to the existence of exact Lagrangian submanifolds and of exact contact hypersurfaces.

Symplectic cohomology is an invariant of symplectic manifolds $(M, d\theta)$ having nice boundaries. For a disc cotangent bundle of a closed manifold, $SH^*(DT^*N)$ is the homology of the free loop space $\mathcal{L}N = C^\infty(S^1, N)$, and Abbondandolo-Schwarz [2] proved that $SH^*(DT^*N)$ has a "pair-of-pants product" which turns into the Chas-Sullivan loop product of string topology [9] on $H_*(\mathcal{L}N)$.

In his survey [33], Seidel mentions that it is possible to define the pair-of-pants product for any M, and more generally TQFT operations

$$\psi_S : SH^*(M)^{\otimes q} \to SH^*(M)^{\otimes p} \qquad (p \geq 1, q \geq 0),$$

by counting maps $S \to M$ from punctured Riemann surfaces S as in Figure 1, satisfying a perturbed Cauchy-Riemann equation, converging to closed 1-periodic Hamiltonian orbits at the punctures.



Figure 1. Pair of pants P; cap C; genus 2 surface S with $(p, q) = (2, 3)$.

For closed symplectic manifolds such a TQFT was constructed on Floer cohomology by Piunikhin-Salamon-Schwarz [26] and Schwarz [32] over fields of characteristic 2. For symplectic manifolds with boundary the construction is more difficult and

---

Date: version: December 2, 2011.

no detailed construction is available in the literature: even the product has only been constructed in detail for cotangent bundles in [1].

Since the aim of this paper will be to construct a deformed TQFT on twisted symplectic cohomology, it has become necessary to fill this gap in the literature. Thus in the Appendix we construct the TQFT in great detail over any field.

Applications arise from proving a vanishing result of $SH^*(M)$, for this reason the most important part of the TQFT is the unit and the product. Indeed $SH^*(M)$ vanishes if and only if the unit vanishes, since then $y = y \cdot 1 = y \cdot 0 = 0$ for any $y \in SH^*(M)$. Secondly, if $W \subset M$ is a nice subdomain then there is a unital ring homomorphism called *Viterbo restriction map*,

$$\varphi : SH^*(M) \to SH^*(W).$$

So if $SH^*(M) = 0$ then $1 = \varphi(1) = \varphi(0) = 0$ in $SH^*(W)$, and so $SH^*(W) = 0$. This circle of ideas is due to Viterbo, Seidel and McLean [36, 33, 22].

0.2. **Deformed operations.** The aim of this paper is to construct a TQFT on the *twisted symplectic cohomology* $SH^*(M)_\alpha$. This is an invariant constructed in [27] associated to a closed 1-form $\alpha \in H^1(\mathcal{L}M)$ on the free loop space, and for $\alpha = 0$ it recovers $SH^*(M)$. In fact it is the Novikov homology theory applied to SH^*, which involves using twisted coefficients in a bundle of formal Laurent series rings.

This twisting yields very concrete applications in symplectic topology, for example in [27] we used it to prove that there are no exact Klein bottles in T^*S^2.

Unfortunately there is no TQFT structure on $SH^*(M)_\alpha$ in general because there is no canonical way of viewing surfaces in M (such as pairs of pants) as chains in $\mathcal{L}M$. Even when a TQFT can be defined on $SH^*(M)_\alpha$, it may not possess a unit for the same reasons that the Novikov cohomology of a manifold usually does not.

Surprisingly, however, we prove that if the form $\alpha \in H^1(\mathcal{L}M)$ is a transgression of a two-form $\eta \in H^2(M)$, then the TQFT structure exists and possesses a unit.

We prove that the restriction maps for twisted symplectic cohomology [27],

$$\varphi : SH^*(M)_\eta \to SH^*(W)_{\eta|_W},$$

are TQFT maps, in particular they are unital ring homomorphisms. Therefore, a vanishing result for $SH^*(M)_\eta$ implies the vanishing of $SH^*(W)_{\eta|_W}$. This entails interesting applications, because by [27] for a simply connected closed manifold N,

$$SH^*(T^*N)_\eta \cong H_{n-*}(\mathcal{L}N)_\eta = 0 \text{ for non-zero } \eta \in H^2(N).$$

0.3. **Topological quantum field theory.** A *TQFT* [4] is a tensor functor from the category of 2-dimensional cobordisms between 1-manifolds to the category of vector spaces. We make this explicit in our setup: to a circle we associate the vector space $SH^*(M)_\eta$, the symplectic cohomology of M computed over a field $\mathbb{K}$ of coefficients (twisted by $\eta \in H^2(M)$, when twisting is present). To the disjoint union of p circles we associate the tensor product $SH^*(M)_\eta^{\otimes p}$ of p copies of $SH^*(M)_\eta$, and to the empty set we associate the base field $\mathbb{K}$. On morphisms, the functor is

$$(\text{Punctured Riemann surface } S) \mapsto (\text{Operation } \psi_S : SH^*(M)_\eta^{\otimes q} \to SH^*(M)_\eta^{\otimes p}),$$

where S represents a cobordism between $p \geq 1$ and $q \geq 0$ circles (Figure 1). Functoriality means that $\psi_Z = \mathrm{id}$ for cylinders $Z : p = q = 1$ and that compositions are respected: gluing surfaces along cylindrical ends yields compositions of ψ_S maps. The TQFT is topological since ψ_S only depends on p, q and the genus of S.

Theorem. *$SH^*(M)_\eta$ is a $(1+1)$-dimensional TQFT, except it does not possess operations for $p = 0$ (so no counit).*

The ψ_S are not defined for $p = 0$ due to a non-compactness issue. Indeed if $p = 0$ were allowed, then $SH^*(M)$ would be finite dimensional, which is false for simply connected cotangent bundles. This differs from Floer cohomology of (suitable) *closed* symplectic M: $p = 0$ is legitimate, $FH^*(M) \cong H^*(M)$ is finite dimensional.

We summarize the untwisted TQFT below (in 2.7 we prove $SH_*(M)$ is canonically the dual of $SH^*(M)$, which had not been clarified in the literature before).

		Symplectic Cohomology	deg	Symplectic Homology	deg
$H =$ Hamiltonian linear at ∞ 		$SC^*(H) = \bigoplus_{\text{1-periodic orbits } x} \mathbb{K}x$ $d : SC^*(H) \to SC^{*+1}(H)$ $dy = \sum_{u \in \mathcal{M}_0(x,y)} \varepsilon_u\, x$	1	$SC_*(H) = \prod_{\text{1-periodic orbits } x} \mathbb{K}x$ $\delta : SC_*(H) \to SC_{*-1}(H)$ $\delta x = \sum_{u \in \mathcal{M}_0(x,y)} \varepsilon_u\, y$	-1
slopes at ∞ : $m \leq m'$ 		Continuation maps $\varphi : SC^*(H^m) \to SC^*(H^{m'})$ $\varphi(y) = \sum_{v \in \mathcal{M}_0^{H_s}(x,y)} \varepsilon_v\, x$	0	Continuation maps $SC_*(H^m) \leftarrow SC_*(H^{m'}) : \varphi$ $\varphi(x) = \sum_{v \in \mathcal{M}_0^{H_s}(x,y)} \varepsilon_v\, y$	0
		$SH^*(M) = \varinjlim SH^*(H)$		$SH_*(M) = \varprojlim SH_*(H)$	
Canonical map		$c^* : H^*(M) \to SH^*(M)$	0	$c_* : SH_*(M) \to H_*(M)$	0
Viterbo restriction $\left(W \subset M\right.$ Liouville $\left.\text{subdomain}\right)$		$SH^*(M) \to SH^*(W)$	0	$SH_*(W) \to SH_*(M)$	0
String topology		$SH^*(T^*N) \cong H_{n-*}(\mathcal{L}N)$		$SH_*(T^*N) \cong H^{n-*}(\mathcal{L}N)$	
Surface S	(p,q,g)	TQFT structure	deg	TQFT structure	deg
	$(1,0,0)$	Unit $\psi_C : \mathbb{K} \to SH^*(M)$	0	Counit $\psi^C : SH_*(M) \to \mathbb{K}$	0
	$(1,1,0)$	Identity $\psi_Z = \mathrm{id} : SH^*(M) \to SH^*(M)$	0	Identity $\psi^Z = \mathrm{id} : SH_*(M) \to SH_*(M)$	0
	$(1,2,0)$	Product $\psi_P : SH^*(M)^{\otimes 2} \to SH^*(M)$	0	Coproduct $\psi^P : SH_*(M) \to SH_*(M)^{\otimes 2}$	0
	$(2,1,0)$	Coproduct $\psi_Q : SH^*(M) \to SH^*(M)^{\otimes 2}$	$2n$	Product $\psi^Q : SH_*(M)^{\otimes 2} \to SH_*(M)$	$-2n$
	$(3,2,4)$	$\psi_S : SH^*(M)^{\otimes q} \to SH^*(M)^{\otimes p}$ degree $= -2n(1-g-p)$ $p \geq 1,\ q \geq 0$		$\psi^S : SH_*(M)^{\otimes p} \to SH_*(M)^{\otimes q}$ degree $= 2n(1-g-p)$ $p \geq 1,\ q \geq 0$	

Corollary. *There is a graded-commutative associative unital ring structure on $SH^*(M)_\eta$ with product ψ_P and unit $e = \psi_C(1)$ (defined also over integer coefficients).*

0.4. **TQFT structure on ordinary cohomology.** A TQFT structure exists also on the ordinary cohomology $H^*(M)$ if we use 1-dimensional cobordisms. Identify the ordinary cohomology with the Morse cohomology $MH^*(f)$ of any Morse function $f : M \to \mathbb{R}$. Replace surfaces S by directed graphs S'. Consider for example the Y-shaped graph P' in Figure 12. Assign a generic Morse function f_i to each edge. The count of isolated negative gradient flow lines along the graph defines

$$\psi_{P'} : MH^*(f_2) \otimes MH^*(f_3) \to MH^*(f_1).$$

After identifications with ordinary cohomology, this is the cup product on $H^*(M)$. Indeed $H^*(M)$ is a $(0+1)$-dimensional TQFT: to a point associate $H^*(M)$, and to a cobordism between $p+q$ points represented by a directed graph S' associate

$$\psi_{S'} : H^*(M)^{\otimes q} \to H^*(M)^{\otimes p}.$$

This TQFT is well-known, see for example Betz-Cohen [5] and Fukaya [19].

0.5. **Wrapped Floer cohomology.** *Wrapped Floer cohomology* $HW^*(L)$ are invariants of an exact Lagrangian $L \subset (M, d\theta)$ having Legendrian intersection with ∂M. Special cases for $M = T^*N$ were introduced by Abbondandolo-Schwarz [1], the general definition arises in Fukaya-Seidel-Smith [20], and a detailed construction is in Abouzaid-Seidel [3]. The construction in [3] is more complicated than ours because the authors' aim was to construct an A_∞-structure at the chain level.

The construction of $SH^*(M)$ involves closed Hamiltonian orbits whereas $HW^*(L)$ involves open Hamiltonian orbits with ends on L. This open-closed string theory analogy is a dictionary to pass from SH^* to HW^*: we now use half the surface obtained after cutting Figure 1 with a vertical plane, so S has boundary components. We count maps $u : S \to M$ satisfying the Lagrangian boundary condition $u(\partial S) \subset L$, converging to open Hamiltonian orbits at the $p+q$ punctures, yielding:

$$\mathcal{W}_S : HW^*(L)^{\otimes q} \to HW^*(L)^{\otimes p}. \qquad (p \geq 1, q \geq 0)$$

One can also define open-closed operations by making r punctures in the interior of S. For a disc $S = D$ with $(p, q, r) = (1, 1, 1)$ this yields a module structure

$$\mathcal{W}_D : SH^*(M) \otimes HW^*(L) \to HW^*(L).$$

This paper will introduce a new group: the *twisted wrapped Floer cohomology*, $HW^*(L)_\eta$, which depends on a closed two-form $\eta \in H^2(M, L)$. This recovers $HW^*(L)$ when $\eta = 0$. Denote $\overline{\eta}$ the image of η under $H^2(M, L) \to H^2(M)$.

Theorem. *$HW^*(L)_\eta$ has a TQFT structure, part of which is a unital ring structure. It is an $SH^*(M)_{\overline{\eta}}$-module via $\mathcal{W}_D$. Moreover, the Viterbo restriction maps $HW^*(L)_\eta \to HW^*(L \cap W)|_{\eta|W}$ preserve the TQFT and module structure.*

Corollary. *If $SH^*(M)_{\overline{\eta}}{=}0$ then $HW^*(L)_\eta{=}0$.*

0.6. **The Arnol'd chord conjecture.** It states that a contact manifold containing a Legendrian submanifold must contain a Reeb chord with ends on the Legendrian. We always deal with the Legendrian ∂L of 0.5:

Theorem. *If $SH^*(M)_\eta = 0$ or $HW^*(L)_\eta = 0$ then the chord conjecture holds, and for a generic contact form there are at least rank $H^*(L)$ chords.*

This applies for example to subcritical Stein manifolds M since $SH^*(M) = 0$ by Cieliebak [10]. In this example, the existence of one Reeb chord for any Legendrian $K \subset \partial M$ is due to Mohnke [25]. We will also deduce that:

Corollary. *If M admits an exact embedding into an exact convex symplectic manifold X, and the boundary ∂M is displaceable in X, then the chord conjecture holds for any ∂L, and generically there are rank $H^*(L)$ chords.*

Theorem. *For $L \subset M = DT^*N$, with N closed and simply connected, such that $H^2(T^*N) \to H^2(L)$ is not injective, the chord conjecture holds and generically there are $\geq$ rank $H^*(L)$ chords. It also holds after attaching subcritical handles to DT^*N.*

An ALE space $\overline{M}$ is a simply connected hyperkähler $4-$manifold which at infinity looks like $\mathbb{C}^2/G$ for a finite subgroup $G \subset SL(2, \mathbb{C})$. It arises [28] by symplectizing nice M, such as the plumbing of copies of T^*S^2 by ADE Dynkin diagrams.

Theorem. *For any ALE space the chord conjecture holds for any ∂L and generically there are at least rank $H^*(L)$ Reeb chords.*

0.7. **Obstructions to exact contact embeddings.** $j : \Sigma^{2n-1} \hookrightarrow (M^{2n}, d\theta)$ is an *exact contact embedding* if there is a contact form α on Σ with $\alpha - j^*\theta =$ exact. For example, if $L \subset M$ is a closed exact Lagrangian, then a Weinstein neighbourhood $W \cong DT^*L$ yields an exact contact hypersurface $ST^*L \cong \partial W \hookrightarrow M$. Using the deformed TQFT we prove the theorem: (stronger results are discussed in 11.4, 11.5)

Theorem. *Let L, N be closed simply connected n-manifolds, $n \geq 4$. For any exact contact embedding $ST^*L \hookrightarrow T^*N$, the following hold*

1. *$H^2(N) \to H^2(L)$ is injective;*
2. *$\pi_2(L) \to \pi_2(N)$ has finite cokernel;*
3. *if $H^2(N) \neq 0$ then $H_*(L) \cong H_*(W)$ for the filling W of $ST^*L \subset T^*N$.*

0.8. **Displaceability of contact hypersurfaces.** By exploiting recent literature on the Rabinowitz-Floer cohomology $RFH^*(M)$ [11, 12], and after proving that $RFH^*(M) = 0$ iff $SH^*(M) = 0$, we will deduce:

Corollary. *If ∂M is displaceable in the symplectization of M then $SH^*(M) = 0$ and there are no closed exact Lagrangians in M. It also holds if M can be embedded into an exact convex symplectic manifold X, and ∂M is displaceable in X.*

0.9. **String topology.** On the homology of the space ΩN of based loops in N there is the *Pontrjagin product*, induced by concatenating loops to form figure-8 loops. On the homology of the space $\mathcal{L}N$ of free loops in N, there is the *Chas-Sullivan loop product* [9]: given two families of loops, you form the family of all possible figure-8 loops obtained when the two base-points happen to coincide.

Abbondandolo-Schwarz [1] proved that the products on $HW^*(T_q^*N \subset T^*N)$ and $SH^*(T^*N)$ agree with the Pontrjagin and Chas-Sullivan products on $H_{n-*}(\Omega N)$ and $H_{n-*}(\mathcal{L}N)$ via the respective isomorphisms. We extend the result to:

Theorem. *$SH^*(T^*N)_\eta \cong H_{n-*}(\mathcal{L}N)_\eta$ and $HW^*(T_q^*N \subset T^*N)_\eta \cong H_{n-*}(\Omega N)_\eta$ respect the TQFT structures, and the units are $[N]$ and $[$base-point$]$ respectively.*

Acknowledgement: I thank Alberto Abbondandolo, Mohammed Abouzaid, Gabriel Paternain, Matthias Schwarz, and Ivan Smith for many stimulating conversations.

1. Liouville domains

1.1. **Definition.** A Liouville domain $(M^{2n}, \omega = d\theta)$ is a compact exact symplectic manifold with boundary, such that the Liouville vector field Z defined by $\omega(Z, \cdot) = \theta$ points strictly outwards along the boundary ∂M. The form $\alpha = \theta|_{\partial M}$ is a contact form on ∂M, and it defines the Reeb vector field $\mathcal{R}$ on ∂M by the conditions

$$\alpha(\mathcal{R}) = 1,\ d\alpha(\mathcal{R}, \cdot) = 0.$$

The closed orbits of $\mathcal{R}$ on ∂M are called *Reeb orbits*. For a generic choice of α (e.g. a generic choice of θ subject to $d\theta = \omega$) the periods of the Reeb orbits form a countable closed subset of $[0, \infty)$, which we will assume.

The symplectization $\overline{M}$ of M is obtained by gluing a symplectic collar onto ∂M:

$$\overline{M} = M \cup_{\partial M} [0, \infty) \times \partial M.$$

The flow of Z for time $r \leq 0$ parametrizes a neighbourhood $(-\varepsilon, 0] \times \partial M$ of ∂M. All data then naturally extends to $\overline{M}$: $Z = \partial_r$, $\theta = e^r\alpha$, $\omega = d\theta$.

Example. The disc bundle $M = DT^*N$ of a closed Riemannian manifold N^n, in local coordinates (q_i, p_i): $\theta = \sum p_i\, dq_i$, $\omega = \sum dp_i \wedge dq_i$, $Z = \sum p_i \frac{\partial}{\partial p_i}$, $\overline{M} = T^*N$.

The flow of Z for time $r \in (-\infty, \infty)$ defines the coordinate $R = e^r \in (0, \infty)$ on $\overline{M}$. We will use R instead of r from now on, so $\overline{M} = M \cup [1, \infty) \times \partial M$.

Let J be an ω-compatible almost complex structure on $\overline{M}$. Let $g = \omega(\cdot, J\cdot)$ denote the J-invariant metric. We always assume that J is of *contact type*:

$$J^*\theta = dR \quad \text{(for large } R\text{)}.$$

For $H \in C^\infty(\overline{M}, \mathbb{R})$, define the *Hamiltonian vector field* $X = X_H$ by $\omega(\cdot, X) = dH$.

We always assume H is C^2-small and Morse inside M. So the only 1-periodic orbits inside M are constants: the critical points of H. We also assume that $H = h(R)$ depends only on R on the collar. So $X = h'(R)\mathcal{R}$, so 1-periodic orbits of X on the collar have constant $R = R(x)$ and they correspond to the Reeb orbits y of period $T = h'(R)$ via $y(t) = x(t/T)$. We always assume h is linear for large R with generic slope, so no 1-periodic Hamiltonian orbits appear there.

1.2. **Action functional.** Let $\mathcal{L}\overline{M} = C^\infty(S^1, \overline{M})$ be the space of free loops in $\overline{M}$. Let A_H denote the H-perturbed action functional for $x \in \mathcal{L}\overline{M}$,

$$A_H(x) = -\int x^*\theta + \int_0^1 H(x(t))\, dt.$$

The differential of A_H at $x \in \mathcal{L}\overline{M}$ in the direction $\xi \in T_x\mathcal{L}\overline{M} = C^\infty(S^1, x^*T\overline{M})$ is $dA_H \cdot \xi = -\int_0^1 \omega(\xi, \dot{x} - X)\, dt$, so the critical points of A_H are the 1-periodic orbits $\dot{x}(t) = X(x(t))$. For an orbit x on the collar in $\{R\} \times \partial M$, $A_H(x) = -Rh'(R) + h(R)$.

1.3. **Floer trajectories.** With respect to the L^2-metric $\int_0^1 g(\cdot,\cdot)\, dt$ the gradient is $\nabla A_H = J(\dot{x} - X)$. For $u : \mathbb{R} \to \mathcal{L}\overline{M}$, or equivalently $u : \mathbb{R} \times S^1 \to \overline{M}$, the negative L^2-gradient flow equation $\partial_s u = -\nabla A_H(u)$ in the coordinates $(s, t) \in \mathbb{R} \times S^1$ is

$$\partial_s u + J(\partial_t u - X) = 0 \quad \text{(Floer's equation)}.$$

Let $\widehat{\mathcal{M}}(x_-, x_+)$ denote the solutions u converging to 1-periodic Hamiltonian orbits $x_\pm$ at the ends $s \to \pm\infty$. Then $\mathcal{M}(x_-, x_+) = \widehat{\mathcal{M}}(x_-, x_+)/\mathbb{R}$ denotes the moduli space of *Floer trajectories*, where we have identified $u(\cdot, \cdot) \sim u(\cdot + \text{constant}, \cdot)$.

1.4. **Energy.** The energy of a Floer trajectory $u \in \mathcal{M}(x_-, x_+)$ is defined as

$$\begin{aligned} E(u) &= \int |\partial_s u|^2\, ds \wedge dt \\ &= \int d\theta(\partial_s u, \partial_t u - X)\, ds \wedge dt \\ &= \int u^* d\theta - dH(\partial_s u)\, ds \wedge dt \\ &= \int_0^1 (\theta(\dot{x}_+) - \theta(\dot{x}_-) - H(x_+) + H(x_-))\, dt \\ &= A_H(x_-) - A_H(x_+). \end{aligned}$$

Thus we have an a priori energy estimate (in terms only of the ends $x_\pm$ not of u).

1.5. **Transversality and compactness.** Standard Floer theory methods [29] show that for a generic time-dependent perturbation (H_t, J_t) of (H, J) the 1-periodic Hamiltonian orbits are isolated and the moduli spaces $\mathcal{M}(x_-, x_+)$ are smooth manifolds. The time-dependence of H_t actually only needs to be switched on near the ends, so as to guarantee that the Hamiltonian orbits are isolated (and nondegenerate). Write $\mathcal{M}_k(x_-, x_+)$ for the k-dimensional part of $\mathcal{M}(x_-, x_+)$.

Lemma 1 (Consequence of Lemma 81)**.** *Solutions of* $\partial_s u + J(\partial_t u - X) = 0$ *converging to* $x_\pm$ *at the ends are entirely contained in the region* $R \leq \max R(x_\pm)$.

By this maximum principle, the Floer trajectories in $\mathcal{M}(x_-, x_+)$ stay in a compact region of $\overline{M}$. Thus we reduce to checking whether the compactness proofs that hold for closed symplectic manifolds (e.g. Salamon [29]) are applicable. Indeed, we have the two sufficient requirements: an a priori energy estimate and a reason to exclude the bubbling-off of J-holomorphic spheres (there are no non-constant J-holomorphic spheres by Stokes's theorem since $\omega = d\theta$ is exact).

Therefore, the moduli spaces $\mathcal{M}(x_-, x_+)$ have natural compactifications, whose boundaries are described by broken Floer trajectories (see Figure 8). In particular $\mathcal{M}_0(x_-, x_+)$ is already compact, so it is a finite set of points (*isolated solutions*).

Convention. *We write* (H, J) *even though one actually uses a perturbed* (H_t, J_t).

2. Symplectic cohomology and symplectic homology

2.1. **Symplectic chain complex.** Pick a base field $\mathbb{K}$. Let $SC^*(H)$ denote the $\mathbb{K}$-vector space generated by the 1-periodic orbits of X,

$$SC^*(H) = \bigoplus \left\{ \mathbb{K}\, x : x \in \mathcal{L}\overline{M},\ \dot{x}(t) = X(x(t)) \right\}.$$

The differential d is defined on generators by counting isolated Floer trajectories,

$$dy = \sum_{u \in \mathcal{M}_0(x,y)} \epsilon_u\, x$$

where $\epsilon_u \in \{\pm 1\}$ are orientation signs (see Appendix 16), and extend d linearly. By convention, the constant solution $u(s,t) = y(t) \in \mathcal{M}_0(y,y)$ is not counted.

A standard argument [29] shows that $d \circ d = 0$. When gradings are defined (2.6), $\dim \mathcal{M}(x,y) = |x| - |y| - 1$, so we require $|x| - |y| = 1$ in the above definition.

2.2. **Continuation maps.** Let H_s, J_s depend on s, relative to the ends ($H_s = H_\pm$, $J_s = J_\pm$ for large $|s|$). The moduli space $\mathcal{M}^{H_s}(x_-, x_+)$ of *Floer continuation solutions* consists of the solutions $v : \mathbb{R} \times S^1 \to \overline{M}$ of $\partial_s v + J_s(\partial_t v - X_{H_s}) = 0$, which converge to $X_{H_\pm}$-orbits $x_\pm$ as $s \to \pm\infty$ (without making any identifications of solutions). The maximum principle (Lemma 81) holds for *monotone homotopies*:

$$\text{for large } R \text{ we assume: } H_s = h_s(R),\ \partial_s h_s' \leq 0,\ J_s \text{ of contact type.}$$

Say this holds for $R \geq R_0$, then all $v \in \mathcal{M}^{H_s}(x_-, x_+)$ are contained in the compact C defined by $R \leq \max(R(x_-), R(x_+), R_0)$. Suppose that H_s, J_s are s-dependent only for $a \leq s \leq b$, then we get an a priori energy estimate for all $v \in \mathcal{M}^{H_s}(x_-, x_+)$,

$$\begin{array}{rcl} E(u) & = & \int |\partial_s v|^2_{g_s}\, ds \wedge dt \\ & = & A_{H_-}(x_-) - A_{H_+}(x_+) + \int (\partial_s H_s)(v)\, ds \wedge dt \\ & \leq & A_{H_-}(x_-) - A_{H_+}(x_+) + (b-a) \cdot \sup_{m \in C} |\partial_s H_s(m)|. \end{array}$$

Therefore for generic J_s and after a C^2-small time-dependent perturbation, the $\mathcal{M}^{H_s}(x_-, x_+)$ are smooth manifolds with compactifications by broken trajectories.

For monotone H_s, counting isolated Floer continuation solutions defines a *continuation map* $\varphi : SC^*(H_+) \to SC^*(H_-)$. On generators

$$\varphi(x_+) = \sum_{v \in \mathcal{M}_0^{H_s}(x_-, x_+)} \epsilon_v\, x_-$$

where $\epsilon_v \in \{\pm 1\}$ are orientation signs (see Appendix 16). Then extend φ linearly. A standard argument [29] shows that φ is a chain map. When gradings are defined, $\dim \mathcal{M}^{H_s}(x_-, x_+) = |x_-| - |x_+|$, so we require $|x_-| = |x_+|$ in the above definition.

Lemma 2 (for example [27])**.** *Changing the monotone homotopy (H_s, J_s) changes φ by a chain homotopy. So on cohomology $\varphi : SH^*(H_+) \to SH^*(H_-)$ is independent of (H_s, J_s). Also φ equals any composite $SH^*(H_+) \to SH^*(K) \to SH^*(H_-)$ induced by the continuation maps for monotone homotopies H_- to K and K to H_+. The constant homotopy $H_s = H$ induces the identity on $SH^*(H)$. If $H_\pm$ are linear at infinity with the same slope then φ is an isomorphism on cohomology.*

2.3. **Hamiltonians linear at infinity.** Let H^m be any Hamiltonian equal to $mR+$constant for large R (generic $m > 0$). By Lemma 2, $SH^*(H^m)$ only depends on m and continuation maps $SH^*(H^m) \to SH^*(H^{m'})$ exist for $m \leq m'$.

2.4. **Symplectic cohomology.** Define the *symplectic cohomology* by

$$SH^*(M) = SH^*(\overline{M}) = \varinjlim SH^*(H),$$

where the direct limit is taken over the continuation maps between Hamiltonians linear at infinity. In particular, $SH^*(M) \cong \varinjlim SH^*(H^m)$ for any sequence $m \to \infty$.

In Appendix 17 we mention the construction of $SH^*(M)$ using just one Hamiltonian which is non-linear at infinity. This makes many arguments unnecessarily complicated (see 17.2) so for now we assume our Hamiltonians are linear at infinity.

2.5. **Invariance under symplectomorphisms of contact type.** Let M, N be Liouville domains. A symplectomorphism $\varphi : \overline{M} \to \overline{N}$ is of *contact type* if

$$\varphi^*\theta_N = \theta_M + d(\text{compactly supported function}).$$

It follows that at infinity $\varphi(e^r, y) = (e^{r-f(y)}, \psi(y))$, where $y \in \partial M$, for a smooth $f : \partial M \to \mathbb{R}$ and a contact isomorphism $\psi : \partial M \to \partial N$ with $\psi^*\alpha_N = e^f \alpha_M$.

Under such a map $\varphi : \overline{M} \to \overline{N}$, the Floer solutions on $\overline{N}$ for $(H, d\theta_N, J_N)$ corresponds precisely to the Floer solutions on $\overline{M}$ for $(\varphi^*H, d\theta_M, \varphi^*J_N)$. However, for H on $\overline{N}$ linear at infinity, $\varphi^*H(e^r, y) = h(e^{r-f(y)})$ is not linear at infinity.

So $SH^*(N)$ is isomorphic via φ^* to $SH^*_f(M) = \varinjlim SH^*(H_f)$ calculated for Hamiltonians of the form $H_f = h(R_f)$, with $R_f = e^{r-f(y)}$, using almost complex structures J satisfying $J^*d\theta = dR_f$. It still turns out that $SH^*(M) \cong SH^*_f(M)$. This is proved by a continuation argument by homotoping f to zero and proving a maximum principle for $R_f \circ u$ for such homotopies (e.g. see [28, Lemma 7]).

Lemma 3. *A contact type $\varphi : \overline{M} \cong \overline{N}$ induces $\varphi_* : SH^*(M) \cong SH^*(N)$.*

2.6. **Grading of symplectic cohomology.** When we use a $\mathbb{Z}$-grading on $SH^*(M)$ we assume that $c_1(M) = 0$, which ensures the following index $|\cdot|$ is defined.

Since $c_1(M) = 0$, we can pick a trivialization of the canonical bundle $\mathcal{K} = \Lambda^{n,0}T^*M$. Over any 1-periodic Hamiltonian orbit x, trivialize x^*TM so that it induces an isomorphic trivialization of $\mathcal{K}$. Let ϕ_t denote the linearization $D\varphi^t(\gamma(0))$ of the time t Hamiltonian flow written in a trivializing frame for x^*TM.

Let $\operatorname{sign}(t)$ denote the signature of the quadratic form $\omega(\cdot, \partial_t\phi_t \cdot) : \ker(\phi_t - \mathrm{id}) \to \mathbb{R}$, assuming we perturbed ϕ_t relative endpoints to make the quadratic form nondegenerate and to make $\ker(\phi_t - \mathrm{id}) = 0$ except at finitely many t.

The *Maslov index* $\mu(\gamma)$ of γ is $\mu(\gamma) = \frac{1}{2}\operatorname{sign}(0) + \sum_{0<t<1} \operatorname{sign}(t) + \frac{1}{2}\operatorname{sign}(1)$.

The Maslov index is invariant under homotopy relative endpoints, and it is additive with respect to concatenations. If ϕ_t is a loop of unitary transformations, then its Maslov index is the winding number of the determinant, $\det \phi_t : \mathcal{K} \to \mathcal{K}$. For example $\phi_t = e^{2\pi i t} \in U(1)$ for $t \in [0,1]$ has Maslov index 1.

The $\mathbb{Z}$-grading on $SH^*(M)$ when $c_1(M) = 0$ is the *Conley-Zehnder index*:

$$|x| = \tfrac{\dim(M)}{2} - \mu(x).$$

When $c_1(M) \neq 0$ this still defines a $\mathbb{Z}/2\mathbb{Z}$ grading since the Conley-Zehnder indices will change by an integer multiple of $2c_1(M)$ when we change the trivialization over a Hamiltonian orbit. In particular, Koszul signs $(-1)^{|x|}$ are defined (see 16.15). We refer to Salamon [29] for a detailed exposition on gradings.

Remark 4. *Our convention ensures that $|x|$ equals the Morse index of x when x is a critical point of a C^2-small Morse Hamiltonian H. This ensures that the unit of $SH^*(M)$ lies in degree 0, and the product is additive on degrees. This differs from Schwarz's convention* [32], *$\|x\| = n - ind_{Morse}(x)$, which explains why the index of Lemma 83 is $-\sum \|x_a\| + \sum \|y_b\| + n \cdot \chi(S)$ in* [32], *with $\chi(S) = 2 - 2g - p - q$.*

2.7. **Symplectic homology.** In the notation of 2.1, define:

$$SC_*(H) = \prod \left\{ \mathbb{Z}x : x \in \mathcal{L}\overline{M},\ \dot{x}(t) = X_H(x(t)) \right\}$$

$$\delta : SC_*(H) \to SC_{*-1}(H),\ \ \delta x = \sum_{u \in \mathcal{M}_0(x,y)} \epsilon_u\, y \quad \text{(compare 2.1)}$$

$$\varphi_* : SC_*(H_-) \to SC_*(H_+),\ \ \varphi_*(x_-) = \sum_{v \in \mathcal{M}_0^{H_s}(x_-,x_+)} \varepsilon_v\, x_+ \quad \text{(compare 2.2)}$$

$$SH_*(M) = \varprojlim SH_*(H) \quad \text{(compare 2.4)}.$$

Notice we count solutions flowing *out* of a generator not *into*. This reverses all maps, so the symplectic homology is the *inverse* limit over continuation maps.

2.8. **Symplectic homology is the dual of symplectic cohomology.** This section aims to clarify once and for all the relation between *symplectic homology* and *symplectic cohomology*, a distinction in terminology which so far had been used in the literature only to distinguish various conflicting sign conventions.

Observe that $SC_*(H) = \mathrm{Hom}(SC^*(H), \mathbb{K})$ canonically and that the differentials δ, d are dual to each other, indeed on generators: $\delta x = \sum A_{x,y}\, y$ and $dy = \sum A_{x,y}\, x$ where $A_{x,y} = \#\mathcal{M}_0(x,y)$ (count the elements u with signs ϵ_u). Thus δ, d are represented by the matrices $(A_{y,x}), (A_{x,y})$, which are transpose to each other. Thus $SC_*(H)$ is canonically the dual of $SC^*(H)$, and so by universal coefficients $SH_*(H)$ is the dual of $SH^*(H)$ and comes with a canonical isomorphism

$$SH_*(H) \to \mathrm{Hom}(SH^*(H), \mathbb{K}).$$

Similarly, $\varphi_* : SC_*(H_-) \to SC_*(H_+)$ is dual to $\varphi^* : SC^*(H_+) \to SC^*(H_-)$ (2.2).

Theorem 5. *Symplectic homology is canonically dual to symplectic cohomology:*

$$SH_*(M) \cong SH^*(M)^\vee.$$

Proof. Using the above canonical isomorphisms, we get the commutative diagram

$$\begin{tikzcd}
SH_*(H_+) \arrow[r, "\sim"] & \mathrm{Hom}(SH^*(H_+), \mathbb{K}) \\
SH_*(H_-) \arrow[r, "\sim"] \arrow[u, "\varphi_*"] & \mathrm{Hom}(SH^*(H_-), \mathbb{K}) \arrow[u, "(\varphi^*)^\vee"']
\end{tikzcd}$$

By category theory, $\mathrm{Hom}(\varinjlim M_i, N) \cong \varprojlim \mathrm{Hom}(M_i, N)$, for any module N and any directed system of modules M_i. Using $SH^*(H), \mathbb{K}$ in place of M_i, N, we deduce

$$SH^*(M)^\vee \equiv \mathrm{Hom}(SH^*(M), \mathbb{K}) \cong \varprojlim \mathrm{Hom}(SH^*(H), \mathbb{K}) \cong \varprojlim SH_*(H) = SH_*(M)$$

using the commutative diagram to obtain the third identification. □

Corollary 6. *Symplectic cohomology is* 0 *if and only if symplectic homology is* 0*.*

3. Wrapped Floer cohomology

3.1. **Lagrangians inside Liouville domains.** Let $(M^{2n}, d\theta)$ be a Liouville domain. Let $L^n \subset M$ be an exact Lagrangian submanifold with Legendrian boundary $\partial L = L \cap \partial M$ such that this intersection is transverse. *Exact Lagrangian* means the pull-back $\theta|_L = \text{exact}$. *Legendrian* means $T\partial L \subset \ker \alpha$, or equivalently $\theta|_{\partial L} = 0$.

We strengthen the last condition: the pull-back $\theta|_L = df$ vanishes near ∂L. This stronger condition can always be achieved for the new data obtained after deforming L by a Hamiltonian isotopy of M relative to ∂M (see [3, Lemma 4.1]).

This condition ensures that near ∂M, L has the form (interval) $\times\ \partial L$ in the coordinates $(0, \infty) \times \partial M \subset \overline{M}$ of 1.1. We extend L to the non-compact Lagrangian

$$\overline{L} = L \cup ([1, \infty) \times \partial L) \subset \overline{M},$$

with $\theta|_{\overline{L}} = df$, and f locally constant on $\overline{L} \setminus L$ since $\theta|_{\overline{L} \setminus L} = 0$.

Example 7. *The fibre* $\overline{L} = T_q^* N$ *inside* $\overline{M} = T^*N$ *for a closed manifold* N*, where* $M = DT^*N$*. More generally, the conormal bundle of a proper submanifold* $S \subset N$*:*

$$\overline{L} = N^*S = \{(q, p) : q \in S, p|_{T_qS} = 0\} \subset T^*N.$$

3.2. **Hamiltonian and Reeb chords.** A *Hamiltonian chord* is a map

$$x : [0, 1] \to \overline{M}, \text{ with } \dot{x}(t) = X_H(x(t)) \text{ and ends } x(0), x(1) \in \overline{L}.$$

A *Reeb chord* of period T is a map

$$y : [0, T] \to \partial M, \text{ with } \dot{y}(t) = \mathcal{R}(y(t)) \text{ and ends } y(0), y(1) \in \partial L.$$

Let φ_H^t denote the time t flow of $X = X_H$. The Hamiltonian chords correspond to the intersections $\varphi_H^1(L) \cap L$. We choose H such that on $R \leq 1$, H is C^2-small and Morse, and on $R \geq 1$, $H = h(R)$, $h' > 0$, and $h' = m$ for $R \gg 0$. It follows that:

Lemma 8. *The Hamiltonian chords in the collar correspond to Reeb chords* y *of period* $T \leq m$ *via* $x \leftrightarrow (y, R)$*, where* $h(R) = H(x)$*,* $y(t) = x(t/T)$*,* $T = h'(R)$*.*

Lemma 9. *For generic* $m, H, \overline{L}$*, there are only finitely many Hamiltonian chords.*

Proof. Unlike the $SH^*(H)$ construction, we will not need to make time-dependent perturbations of H. However, we need to allow perturbations $\psi(L)$ of L by a compactly supported Hamiltonian flow ψ (one can alternatively view this as keeping L fixed but perturbing H to $H \circ \psi^{-1}$, at the cost of losing Lemma 8). For generic $m > 0$ there are no Reeb chords of period m (by a Sard's Lemma argument). After a small generic (time-independent) compactly supported Hamiltonian perturbation of $H, \overline{L}$ one can ensure the Hamiltonian chords are non-degenerate, that is $\varphi_H^1(\overline{L})$ is transverse to $\overline{L}$ ([3, Lemma 8.1]). Thus the Hamiltonian chords are isolated. □

3.3. **Action functional.** Consider the space of smooth paths with ends in $\overline{L}$:

$$\Omega(\overline{M},\overline{L}) = \{x \in \mathbb{C}^\infty([0,1],\overline{M}) : x(0), x(1) \in \overline{L}\}.$$

Define $A_H : \Omega(\overline{M},\overline{L}) \to \mathbb{R}$ analogously to 1.2,

$$A_H(x) = f(x(1)) - f(x(0)) - \int x^*\theta + \int_0^1 H(x(t))\, dt.$$

The motivation for the first three terms is that they would arise by Stokes's theorem from $\int u^*\omega$ if u were a disc with boundary in $\overline{L} \cup \text{image}(x)$. This ensures that

$$dA_H \cdot \xi = -\textstyle\int_0^1 \omega(\xi, \dot{x} - X)\, dt,$$

where $\xi \in T_x\Omega(\overline{M},\overline{L}) = \{\xi \in C^\infty([0,1], x^*T\overline{M}) : \xi(0), \xi(1) \in T\overline{L}\}$.

Therefore the critical points of A_H are the Hamiltonian chords. For Hamiltonian chords x on the collar, $A_H(x) = f(x(1)) - f(x(0)) - Rh'(R) + h(R)$.

3.4. **Wrapped trajectories.** Pick J, g as in 1.1. The solutions $u : \mathbb{R} \to \Omega(\overline{M},\overline{L})$ of $\partial_s u = -\nabla A_H$ are the solutions $u : \mathbb{R} \times [0,1] \to \overline{M}$ of $\partial_s u + J(\partial_t u - X) = 0$, with Lagrangian boundary conditions $u(\cdot,0), u(\cdot,1) \in \overline{L}$. Let $\widehat{\mathcal{W}}(x_-,x_+)$ denote the solutions converging to $x_\pm$ at the ends $s \to \pm\infty$. Let $\mathcal{W}(x_-,x_+) = \widehat{\mathcal{W}}(x_-,x_+)/\mathbb{R}$ denote the moduli space of *wrapped trajectories*, identifying $u(\cdot,\cdot) \sim u(\cdot + \text{constant}, \cdot)$.

Remark 10. *These are the same moduli spaces one would study to define the Lagrangian Floer homology* $HF^*(\varphi_H^1(\overline{L}),\overline{L})$. *Indeed wrapped trajectories* $u(s,t)$ *correspond precisely to pseudo-holomorphic strips with boundaries in* $\varphi_H^1(\overline{L})$ *and* $\overline{L}$, *that is solutions* $v : \mathbb{R} \times [0,1] \to \overline{M}$ *of* $\partial_s v + \tilde{J}_t \partial_t v = 0$, *converging to* $v(\pm\infty,\cdot) = x_\pm \in \varphi_H^1(\overline{L}) \cap \overline{L}$, *where* $\tilde{J}_t = d\varphi_H^{1-t} \circ J \circ d\varphi_H^{t-1}$. ***Proof****. Let* $v(s,t) = \varphi_H^{1-t}(u(s,t))$. $\square$

3.5. **Energy.** The energy of $u \in \mathcal{W}(x_-,x_+)$ is defined as $E(u) = \int |\partial_s u|^2\, ds \wedge dt$. The same calculation as in 1.4, using $\theta|_{\overline{L}} = df$, yields $E(u) = A_H(x_-) - A_H(x_+)$.

3.6. **Transversality and compactness.** Just as in the Floer case, a generic time-dependent perturbation of J ensures that $\mathcal{W}(x_-,x_+)$ are smooth manifolds.

Lemma 11 (Maximum principle, see 15.22)**.** *All wrapped trajectories in* $\mathcal{W}(x_-,x_+)$ *land entirely in the region* $R \leq \max(R(x_-), R(x_+))$.

This and the energy estimate 3.5 imply that $\mathcal{W}(x_-,x_+)$ have compactifications by broken wrapped trajectories (no non-constant J-holomorphic spheres or discs with boundary in $\overline{L}$ can bubble off since $d\theta$, $\theta|_{\overline{L}}$ are exact). Indeed the analysis of $\mathcal{W}(x_-,x_+)$ reduces to the known case of Lagrangian Floer homology (Remark 10).

3.7. **Wrapped Floer cohomology.** Pick a base field $\mathbb{K}$. The wrapped Floer complex $CW^*(H;L)$ is the $\mathbb{K}$-vector space generated by the Hamiltonian chords,

$$CW^*(H;L) = \bigoplus \left\{\mathbb{K}\, x : x \in \Omega(\overline{M},\overline{L}),\ \dot{x}(t) = X_H(x(t))\right\},$$

whose differential d on generators counts isolated wrapped trajectories,

$$dy = \sum_{u \in \mathcal{W}_0(x,y)} \epsilon_u\, x,$$

where $\epsilon_u \in \{\pm 1\}$ are orientation signs (see [3]). The constant solution $u(s,t) = y(t) \in \mathcal{W}_0(y,y)$ is not counted. That $d^2 = 0$ is a standard consequence of 3.6. We call $HW^*(L;H) = H_*(CW^*(H;L);d)$ the wrapped Floer cohomology for H.

Remark 12. *For* $\mathbb{K} = \mathbb{Z}/2\mathbb{Z}$ *orientation signs are not necessary. In general, if the Stiefel-Whitney class* $w_2(L) = 0 \in H^2(L;\mathbb{Z}/2)$ *and the relative Chern class* $2c_1(M,L) = 0 \in H^2(M,L;\mathbb{Z})$ *then* [34] *orientation signs are defined and there is a grading on Hamiltonian chords such that* $\dim \mathcal{W}(x_-,x_+) = |x_-| - |x_+| - 1$.

Remark 13. *Via Remark 10,* $HW^*(L;H) \cong HF^*(\varphi_H^1(\overline{L}),\overline{L})$. *Because of the non-compactness of* $\overline{M}$, *this heavily depends on* H*: as the slope* $m \to \infty$, $\varphi_H^1(\overline{L})$ *wraps more and more times around* $\overline{L}$ *giving rise to more chain generators (intersections).*

As in 2.2, a monotone homotopy H_s defines a wrapped continuation map $\varphi : HW^*(H_+;L) \to HW^*(H_-;L)$ by counting wrapped continuation solutions. These are solutions $v : \mathbb{R} \times [0,1] \to \overline{M}$ of $\partial_s v + J_s(\partial_t v - X_{H_s}) = 0$ with Lagrangian boundary conditions $v(\cdot,0), v(\cdot,1) \in \overline{L}$, converging to Hamiltonian chords on the ends $s \to \pm\infty$. The a priori energy estimate for wrapped continuation solutions is the same as in 2.2 and the maximum principle holds by 15.22. This implies the smoothness and compactifiability of the moduli spaces and the analogue of Lemma 2. Thus $HW^*(L;H)$ only depends on the slope m of H at infinity, and as in 2.3 for $m \leq m'$ there is a wrapped continuation map $HW^*(L;H^m) \to HW^*(L;H^{m'})$.

Definition. Define the *wrapped Floer cohomology* by

$$HW^*(L) = \varinjlim HW^*(L;H),$$

taking the direct limit over wrapped continuation maps between Hamiltonians linear at infinity (2.3). So $HW^*(L) \cong \varinjlim HW^*(L;H^m)$ for any sequence $m \to \infty$.

4. CANONICAL c^* MAPS FROM ORDINARY COHOMOLOGY

For $\delta > 0$ smaller than all Reeb periods, consider H^δ as in 2.3. If H^δ is Morse and C^2-small on M then $SC^*(H^\delta)$ reduces to the Morse complex for H^δ, generated by $\mathrm{Crit}(H^\delta)$ and whose differential counts $-\nabla H^\delta$ trajectories (see 14.3). The Morse cohomology is isomorphic to the ordinary cohomology, so $SH^*(H^\delta) \cong H^*(M;\mathbb{K})$. Since H^δ is part of the direct limit construction, we automatically obtain a map

$$c^* : H^*(M;\mathbb{K}) \to SH^*(M).$$

We often write $H^*(M)$ instead of $H^*(M;\mathbb{K})$. The analogue for the wrapped case is

$$c^* : H^*(L;\mathbb{K}) \to HW^*(L).$$

This arises from $c^* : HW^*(L;H^\delta) \to HW^*(L)$, where H^δ is as above so that all intersections $\varphi_{H^\delta}^1(\overline{L}) \cap \overline{L}$ lie inside M. By Lemma 11, all wrapped trajectories lie in M, so we reduce to a compact setup with $\omega = d\theta$. By Remark 13, $HW^*(L;H^\delta) \cong HF^*(\varphi_{H^\delta}^1(L), L)$, and by Floer [16] this is isomorphic to the Morse cohomology $MH^*(L;H^\delta)$. The idea is: $\varphi_{H^\delta}^1(L)$ lies in a Weinstein neighbourhood $W \cong DT^*L$ of L so it is the $\mathrm{graph}(dk) \subset DT^*L$ of a function $k \in C^\infty(L)$. The intersections $\varphi_{H^\delta}^1(L) \cap L$ correspond to $\mathrm{Crit}(k)$, and by an implicit function theorem argument, if H^δ is C^2-small in M, the evaluation $u \mapsto u(s,0)$ sets up a bijection between pseudo-holomorphic strips in M with boundary in L and $-\nabla k$ trajectories in L.

5. TQFT STRUCTURE

In the Appendix we carry out the detailed construction of the (untwisted) TQFT structures on $SH^*(M)$ and $HW^*(L)$, which we summarize here.

Let (S, j) be a Riemann surface with $p \geq 1$ negative and $q \geq 0$ positive punctures. The complex structure j is fixed. Fix cylindrical parametrizations $s + it$ near the punctures (the *cylindrical ends*). Pick a one-form β on S with $d\beta \leq 0$ and with $\beta = h_a \, dt$, $\beta = k_b \, dt$ on the ends for some constants $h_a, k_b > 0$ called *weights*.

The moduli space $\mathcal{M}(x_a; y_b; S, \beta)$ of *Floer solutions* consists of smooth maps $u : S \to \overline{M}$, with $du - X \otimes \beta$ holomorphic, converging on the ends to Hamiltonian orbits x_a, y_b for $h_a H$, $k_b H$. Generically $\mathcal{M}(x_a; y_b; S, \beta)$ are smooth manifolds and have compactifications by broken Floer solutions. When gradings are defined,

$$\dim \mathcal{M}(x_a; y_b; S, \beta) = \sum |x_a| - \sum |y_b| + 2n(1 - g - p).$$

Fix (S, j, β) and an $H \in C^\infty(\overline{M}, \mathbb{R})$ as in 2.3. Define $\psi_S : \otimes_{b=1}^q SC^*(k_b H) \to \otimes_{a=1}^p SC^*(h_a H)$ on the generators by counting isolated Floer solutions

$$\psi_S(y_1 \otimes \cdots \otimes y_q) = \sum_{u \in \mathcal{M}_0(x_a; y_b; S, \beta)} \epsilon_u \, x_1 \otimes \cdots \otimes x_p,$$

where $\epsilon_u \in \{\pm 1\}$ are orientation signs (Appendix 16). Then extend ψ_S linearly. The ψ_S are chain maps. On cohomology, $\psi_S : \otimes_{b=1}^q SH^*(k_b H) \to \otimes_{a=1}^p SH^*(h_a H)$ is independent of the choices (β, j, J) relative to the ends. Taking direct limits:

$$\psi_S : SH^*(M)^{\otimes q} \to SH^*(M)^{\otimes p} \qquad (p \geq 1, q \geq 0).$$

So $SH^*(M)$ has unit $\psi_C(1)$, product ψ_P, coproduct ψ_Q, but no counit since $p \geq 1$.

For SH_* all arrows are reversed: $\psi^S : \bigotimes_{a=1}^p SC_*(h_a H) \to \bigotimes_{b=1}^q SC_*(k_b H)$, $\psi^S(x_1 \otimes \cdots \otimes x_p) = \sum \epsilon_u \, y_1 \otimes \cdots \otimes y_q$ over $u \in \mathcal{M}_0(x_a; y_b; S, \beta)$. Take inverse limits:

$$\psi^S : SH_*(M)^{\otimes p} \to SH_*(M)^{\otimes q} \qquad (p \geq 1, q \geq 0).$$

So $SH_*(M)$ has counit ψ^C, product ψ^Q, coproduct ψ^P, but no unit since $p \geq 1$.

In the wrapped case, the surface S has boundary and it is parametrized by strip-like ends near the punctures (15.21). The moduli space $\mathcal{W}(x_a; y_b; S, \beta)$ of *wrapped solutions* consists of smooth maps $u : S \to \overline{M}$ with $u(\partial S) \subset \overline{L}$, with $du - X \otimes \beta$ holomorphic, converging on the ends to Hamiltonian chords x_a, y_b for $h_a H$, $k_b H$.

In the Appendix we prove that we obtain TQFT maps independent of β, j, J:

$$\mathcal{W}_S : HW^*(L)^{\otimes q} \to HW^*(L)^{\otimes p} \qquad (p \geq 1, q \geq 0),$$

and we construct the module structure of $HW^*(L)$ over the ring $SH^*(M)$.

In 15.20 we describe the TQFT on $H^*(M)$ (and dually $H_*(M)$) via Morse theory.

We postpone to section 14 the following result, which is quite tricky to prove. As far as we know this has been neither proved nor observed in the literature.

Theorem 14. *$c^* : H^*(M) \to SH^*(M)$, $c_* : SH_*(M) \to H_*(M)$ respect the TQFT so they are unital ring maps (cup product on $H^*(M)$, intersection product on $H_*(M)$).*

6. Twisted theory

6.1. **Novikov bundles.** The Novikov algebra Λ over $\mathbb{K}$ in the formal variable t is:

$$\Lambda = \{ \textstyle\sum_{j=0}^\infty k_j \, t^{a_j} \text{ for any } k_j \in \mathbb{K}, \, a_j \in \mathbb{R}, \text{ with } \lim\limits_{j \to \infty} a_j = \infty \}.$$

Let α be a smooth real singular cocycle representing a class $a \in H^1(\mathcal{L}M; \mathbb{R})$ of the free loop space $\mathcal{L}M = C^\infty(S^1, M)$. The Novikov bundle $\underline{\Lambda}_\alpha$ is a local system of coefficients on $\mathcal{L}M$: over a loop γ its fibre is a copy Λ_γ of Λ, and the parallel translation over a path u in $\mathcal{L}M$ connecting γ to γ' is the multiplication isomorphism

$$t^{\alpha[u]}\colon\ \Lambda_\gamma \to \Lambda_{\gamma'}.$$

Here $\alpha[\cdot]\colon\ C_1(\mathcal{L}M;\mathbb{R}) \to \mathbb{R}$ denotes evaluation on smooth singular one-chains. Explicitly for a de Rham 1-cocycle α, on a smooth path $u : I = [0,1] \to \mathcal{L}M$,

$$\alpha[u] = \textstyle\int_I u^*\alpha = \int_0^1 \alpha(\partial_s u)\, ds.$$

Changing α to $\alpha + df$ yields a change of basis isomorphism $x \mapsto t^{f(x)}x$ for the local systems, so by abuse of notation we write $\underline{\Lambda}_a$ and $a[u]$ instead of $\underline{\Lambda}_\alpha$ and $\alpha[u]$.

6.2. **Transgressions.** Let $ev : \mathcal{L}M \times S^1 \to \mathcal{L}M$ be the evaluation map. Define

$$\tau = \pi \circ ev^* : H^2(M;\mathbb{R}) \xrightarrow{ev^*} H^2(\mathcal{L}M \times S^1;\mathbb{R}) \xrightarrow{\pi} H^1(\mathcal{L}M;\mathbb{R})\ ,$$

where π is projection to the Künneth summand. This is an isomorphism when M is simply connected. Explicitly, for a de Rham $\eta \in H^2(M;\mathbb{R})$ and a path $u \subset \mathcal{L}M$,

$$\tau\eta[u] = \textstyle\int \eta(\partial_s u, \partial_t u)\, ds \wedge dt.$$

In particular, $\tau(\eta)$ vanishes on time-independent paths in $\mathcal{L}M$, so $c^*\underline{\Lambda}_{\tau\eta}$ is trivial when we pull-back by the inclusion of constant loops $c : M \to \mathcal{L}M$.

More generally, $\tau(\eta)$ evaluated on a Floer solution $u : S \to \overline{M}$ is defined by

$$\tau\eta[u] = \textstyle\int_S u^*\eta.$$

This integral is finite despite the non-compactness of S because of the exponential convergence of u to the asymptotes (Lemma 82). Indeed, on an end, $|\partial_s u| \leq ce^{-\delta|s|}$, $\partial_s u + J(\partial_t u - C \cdot X) = 0$ and $|\eta(u)| \leq C'$ for some constants c, δ, C, C'. Thus,

$$\textstyle\int_{\text{end}} \eta(\partial_s u, \partial_t u)\, ds \wedge dt \leq C' \cdot \int_{\text{end}} |\partial_s u| \cdot |\partial_t u| \leq C' \cdot \int_{\text{end}} ce^{-\delta|s|}(ce^{-\delta|s|} + C|X|) < \infty.$$

6.3. **Twisted symplectic (co)homology.** Now introduce weights corresponding to $\alpha \in H^1(\mathcal{L}M;\mathbb{R}) \cong H^1(\mathcal{L}\overline{M};\mathbb{R})$ in the construction of 2.1 (for details, see [27]):

$$SC^*(H)_\alpha = SC^*(H;\underline{\Lambda}_\alpha) = \bigoplus \big\{\Lambda x : x \in \mathcal{L}\overline{M},\ \dot{x}(t) = X_H(x(t))\big\}$$

$$dy = \sum_{u \in \mathcal{M}_0(x,y)} \epsilon_u\, t^{\alpha[u]}\, x \quad \text{(see 2.1)}$$

$$SH^*(H)_\alpha = SH^*(H;\underline{\Lambda}_\alpha) = H_*(SC^*(H;\underline{\Lambda}_\alpha); d)$$

$$\varphi : SC^*(H_+)_\alpha \to SC^*(H_-)_\alpha,\ \varphi(x_+) = \sum_{v \in \mathcal{M}_0^{H_s}(x_-,x_+)} \epsilon_v\, t^{\alpha[v]}\, x_- \quad \text{(see 2.2).}$$

For $(M, d\theta; \alpha)$, define $SH^*(M)_\alpha = SH^*(M;\underline{\Lambda}_\alpha) = \varinjlim SH^*(H)_\alpha$, taking the direct limit over the twisted continuation maps φ above (compare 2.4).

Inserting the local system of coefficients $c^*\underline{\Lambda}_\alpha$ in $MH^*(H^\delta)$ in section 4 yields $c^* : H^*(M; c^*\underline{\Lambda}_\alpha) \to SH^*(M;\underline{\Lambda}_\alpha)$. For $\alpha = \tau(\eta)$, $c^*\underline{\Lambda}_\alpha$ is trivial (6.2), so

$$c^* : H^*(M) \otimes \Lambda \to SH^*(M;\underline{\Lambda}_{\tau\eta}).$$

We sometimes abbreviate $H^*(M;\underline{\Lambda}_\alpha)$ by $H^*(M)_\alpha$, and $H^*(M) \otimes \Lambda$ by $H^*(M)$.

The construction of $SH_*(M)_\alpha$ is dual to the above, using weights $t^{-\alpha[u]}, t^{-\alpha[v]}$.

6.4. **Twisted TQFT.** From now on, use $\alpha = \tau\eta$ (6.2). In 6.5 we mention other options. Abbreviate $SC^*(H)_\eta = SC^*(H; \underline{\Lambda}_{\tau\eta})$, $SH^*(H)_\eta = SH^*(H; \underline{\Lambda}_{\tau\eta})$. Define

$$\begin{aligned} \psi_S : \bigotimes_{b=1}^{q} SC^*(k_b H)_\eta &\to \bigotimes_{a=1}^{p} SC^*(h_a H)_\eta \\ \psi_S(y_1 \otimes \cdots \otimes y_q) &= \sum_{u \in \mathcal{M}_0(x_a; y_b; S, \beta)} \epsilon_u \, t^{\tau\eta[u]} \, x_1 \otimes \cdots \otimes x_p. \qquad \text{(compare 15.10)} \end{aligned}$$

Theorem 15.

1. *The weights $\tau\eta[u] = \int_S u^*\eta$ are locally constant on $\mathcal{M}(x_a; y_b; S, \beta)$.*
2. *The weights $\tau\eta[u]$ are additive under the breaking of Floer solutions: if u_λ converge to a broken solution $u\#v$ or $v\#u$, then $\tau\eta[u_\lambda] = \tau\eta[v] + \tau\eta[u]$.*
3. *$\tau\eta[\cdot]$ is constant on components of the compactification of $\mathcal{M}(x_a; y_b; S, \beta)$.*

Proof. Suppose $w(\cdot, \cdot, \lambda) = u_\lambda(\cdot, \cdot)$ is a smooth path of solutions in $\mathcal{M}(x_a; y_b; S, \beta)$, $\lambda \in [0, 1]$. Now $d\eta = 0$, since η is a closed form, so (1) follows by Stokes:

$$0 = \int_{S\times[0,1]} w^* d\eta = \int_S u_1^*\eta - \int_S u_0^*\eta = \tau\eta[u_1] - \tau\eta[u_0].$$

To prove (2), consider a smooth family $w(\cdot, \cdot, \lambda) = u_\lambda \in \mathcal{M}(x_a; y_b; S, \beta)$ which is parametrized by $\lambda \in [0, 1)$, such that $u_\lambda \to u\#v$ as $\lambda \to 1$ (the case $v\#u$ is similar). Fix $\epsilon > 0$. Pick a large $c > 0$ so that the restrictions u', v' of u, v to the complement of the neighbourhoods $(-\infty, -c) \times S^1$, $(c, \infty) \times S^1$ of all the ends of u, v satisfy

$$|\tau\eta[u] + \tau\eta[v] - \tau\eta[u'] - \tau\eta[v']| < \epsilon.$$

This is possible by the calculation of 6.2, which shows that the integrals evaluated near the ends are arbitrarily small because u, v converge exponentially fast.

Denote by u'_λ the analogous restriction of u_λ. The domain of u'_λ is compact so, by Lemma 88, u'_λ converges C^2-uniformly to u'. Moreover there exist $s_\lambda \in \mathbb{R}$ such that $v'_\lambda(s, t) = u_\lambda(s + s_\lambda, t)|_{s\in[-c,c]}$ converges C^2-uniformly to v'. So for large c,

$$|\tau\eta[u'] + \tau\eta[v'] - \tau\eta[u'_\lambda] - \tau\eta[v'_\lambda]| < \epsilon.$$

Since $\epsilon > 0$ is arbitrary, to conclude the proof of (2) we show that for large c

$$|\tau\eta[u'_\lambda] + \tau\eta[v'_\lambda] - \tau\eta[u_\lambda]| < \epsilon.$$

By Lemma 82, near the ends $|\partial_s u| < Ce^{-\delta|s|}$ and $|\partial_s v| < Ce^{-\delta|s|}$ for some δ, C. For large c, these hold at $|s| = c$, which are the boundaries of the domains for u', v'.

Thus, on an end of u, say $K = (-\infty, -c] \times S^1$, apply Stokes's theorem to $K \times [\lambda, 1]$ as in (1) to deduce that $| \int_K u_\lambda^*\eta - \int_K u^*\eta| = | \int_{\mu\in[\lambda,1],|s|=c} u_\mu^*\eta|$. This quantity is arbitrarily small for large c, λ by the calculation of 6.2, since $|\partial_s u_\mu|$ is bounded by $Ce^{-\delta c}$ for large μ. Similar inequalities hold for the other ends of u and v.

Finally, we need to consider what happens near the breaking. So on the end where the breaking occurs, define $w_\lambda(s, t) = u_\lambda(s, t)|_{s\in[c,-c+s_\lambda]}$. At the boundaries $s = c$, $s = -c + s_\lambda$ the w_λ converge C^2-uniformly to $u(c, t)$, $v(-c, t)$ respectively, so we can assume that $\partial_s w_\lambda$ is bounded by a constant multiple of $e^{-\delta c}$ for large λ. Apply Stokes's theorem to $\cup_{\mu\in[\lambda,1]}(c, -c + s_\mu) \times S^1 \times \{\mu\}$ as in (1) to deduce:

$$\left| \int w_\lambda^*\eta - \int_{s\geq c} u^*\eta - \int_{s\leq -c} v^*\eta \right| = \left| \int_{\mu\in[\lambda,1],s=c} u_\mu^*\eta - \int_{\mu\in[\lambda,1],s=-c+s_\mu} u_\mu^*\eta \right|$$

which again is bounded by a constant multiple of $e^{-\delta c}$, and so is arbitrarily small for large c. Thus (2) follows, and (3) follows from (1) and (2). $\square$

Theorem 16. *The twisted TQFT map* ψ_S *is a chain map, so it induces a map* $\psi_S : \otimes_b SH^*(k_b H)_\eta \to \otimes_a SH^*(h_a H)_\eta$ *independent of the choice of data* (β, j, J) (*Theorem 89*)*. For cylinders* Z *the* ψ_Z *are twisted continuation maps* (*Example 80*)*. Gluing surfaces corresponds to compositions* (*Theorem 91*)*. The* ψ_S *maps are compatible with the twisted continuation maps* (*Theorem 93*)*. In the direct limit,*

$$\psi_S : SH^*(M)_\eta^{\otimes q} \to SH^*(M)_\eta^{\otimes p} \qquad (p \geq 1, q \geq 0).$$

Proof. The claims involve the same moduli spaces used to prove the untwisted results (Appendix). The fact that we count the moduli spaces with correct weights follows by Theorem 15: broken solutions appearing on the boundaries $\partial\mathcal{M}$ of a connected component $\mathcal{M}$ of a moduli space are counted with equal weights. $\square$

Corollary 17. $SH^*(M)_\eta$ *carries a twisted TQFT structure, in particular it is a unital ring. The map* $c^* : H^*(M) \otimes \Lambda \to SH^*(M)_\eta$ *of 6.3 respects the TQFT, using Morse operations on* $H^*(M)$*. In particular the unit is* $e = c^*(1) \in SH^0(M)_\eta$.

Proof. The proof is analogous to that of Theorems 95 and 14, which is proved via Lemma 79. We only need to insert appropriate weights. In the notation of Definition 75, define the twisted versions of the maps ϕ, ψ on generators as follows:

$$\begin{aligned} &\phi : SC^*(H^\delta)_\eta \to MC^*(f) \otimes \Lambda & &\psi : MC^*(f) \otimes \Lambda \to SC^*(H^\delta)_\eta \\ &\phi(y) = \sum_{\substack{v\#u \in \mathcal{M}_0^\phi(x,y) \\ \mathrm{image}(v\#u) \subset M}} \epsilon_{v\#u}\, t^{\tau\eta[v]+\tau\eta[u]}\, x & &\psi(x) = \sum_{u\#v \in \mathcal{M}_0^\psi(y,x)} \epsilon_{u\#v}\, t^{\tau\eta[u]+\tau\eta[v]}\, y \end{aligned}$$

The weights for Morse solutions v are redundant: $t^{\tau\eta[c\circ v]} = 1$ since $c \circ v$ only depends on one variable (see 6.2 on why $c^*\underline{\Lambda}_{\tau\eta}$ is trivial). However, inserting these trivial weights makes it clear that the proof of Theorem 15 can be applied to the above moduli spaces: the weights are locally constant on $\mathcal{M}^\phi(x, y)$, $\mathcal{M}^\psi(y, x)$. $\square$

Lemma 18. *The TQFT structure is independent of the choice of representative* $\eta \in H^2(M; \mathbb{R})$*: changing* η *to* $\eta + df$ *for* $f \in C^\infty(\overline{M}, \mathbb{R})$ *yields a change of basis isomorphism* $x \mapsto t^{-\int x^* f} x$ *which respects the twisted TQFT operations on* $SH^*(M)_\eta$.

Theorem 19. *A* $\varphi : \overline{M} \cong \overline{N}$ *of contact type induces* $\varphi_* : SH^*(M)_\alpha \cong SH^*(N)_{\varphi_*\alpha}$*. For* $\alpha = \tau\eta$*,* φ_* *respects the twisted TQFT structure:* $\varphi_*^{\otimes p} \circ \psi_{S,M} = \psi_{S,N} \circ \varphi_*^{\otimes q}$.

Proof. In Lemma 3 and Theorem 94 insert weights $t^{\alpha[v]}$ in the definition of the continuation maps defining $\varphi_* : SH^*(M) \cong SH^*(N)$, and apply Theorem 15. $\square$

6.5. **TQFT structure for other twistings.** For general $\alpha \in H^1(\mathcal{L}M)$, it does not seem possible to define the TQFT structure on $SH^*(M)_\alpha$. Naively one might divide S into cylinders, the restrictions of u yield 1-chains in $\mathcal{L}M$, then evaluate α and add up. This depends heavily on choices and will not yield an invariant TQFT. We can construct the twisting in two cases: (1) for $\alpha = \tau\widetilde{\eta}$ for $\widetilde{\eta} \in H^2(\widetilde{M})$ taken from the universal cover $\widetilde{M}$; and (2) for pull-backs $\alpha = ev^*\mu$ of a 1-form $\mu \in H^1(M)$ under evaluation at 0, $ev : \mathcal{L}N \to N$. For (2) we obtain a TQFT with no unit, whereas for (1) we obtain a TQFT with unit after restricting to contractible loops.

6.6. **Twisting by 2-forms from the universal cover.** Let $\widetilde{M}$ be the universal cover of M. Then $\tau : H^2(\widetilde{M}) \to H^1(\mathcal{L}\widetilde{M})$ is an isomorphism, and

$$H^1(\mathcal{L}\widetilde{M}) \cong \mathrm{Hom}(H_1(\mathcal{L}\widetilde{M}),\mathbb{Z}) \cong \mathrm{Hom}(\pi_1(\mathcal{L}\widetilde{M}),\mathbb{Z}) \cong \mathrm{Hom}(\pi_2(M),\mathbb{Z}),$$

(via evaluation of $\tau\widetilde{\eta}$ on 1-chains). For the component $\mathcal{L}_0 M$ of contractible loops:

$$H^1(\mathcal{L}_0 M) \cong \mathrm{Hom}(\pi_1(\mathcal{L}_0 M),\mathbb{Z}) \cong \mathrm{Hom}(\pi_1 M \ltimes \pi_2 M,\mathbb{Z}) \cong \mathrm{Hom}(H_1 M \times \pi_2 M,\mathbb{Z}).$$

Via these, $H^1(\mathcal{L}\widetilde{M}) \subset H^1(\mathcal{L}_0 M)$, so we can define $SH^*(M)_{\widetilde{\eta}} = SH^*(M;\underline{\Lambda}_{\tau\widetilde{\eta}})$, which comes with $c^* : H^*(M)\otimes\Lambda \to SH^*_0(M)_{\widetilde{\eta}}$. Restricting to $\mathcal{L}_0 M$ as in 15.13:

Theorem 20. *$SH^*_0(M)_{\widetilde{\eta}}$ is a TQFT with unit $\psi_C(1) \in SH^0_0(M)_{\widetilde{\eta}}$ and c^* respects the TQFT. The analogues of Theorems 32, 35, 68 hold.*

Proof. Let α denote a singular 1-cocycle representing the image of $\tau\widetilde{\eta}$ in $H^1(\mathcal{L}_0 M)$. Pick as base point for $\mathcal{L}_0 M$ the constant loop m at a point in M.

Let $u : S \to \overline{M}$ be a Floer solution whose ends x_i are contractible Hamiltonian orbits. Pick maps $\mathbb{C} \to \overline{M}$, whose images are caps C_i which contract x_i down to m. An appropriate gluing $\sigma = (\cup C_i)\#u(S)$ defines a sphere $\sigma : S^2 \to \overline{M}$. Now α corresponds to a homomorphism $\pi_2(M) \to \mathbb{Z}$ under the identification $H^1(\mathcal{L}_0 M) \cong \mathrm{Hom}(\pi_1 M \ltimes \pi_2 M,\mathbb{Z})$ (α vanishes on $\pi_1(M)$). Thus $\alpha[\sigma]$ is defined (using that $\mathcal{L}_0 M \simeq \mathcal{L}_0\overline{M}$ are homotopy equivalent).

Parametrizing $\mathbb{C}$ by polar coordinates, the caps C_i can be viewed as paths in $\mathcal{L}_0 M$ from m to x_i. So C_i is a singular 1-chain in $\mathcal{L}_0 M$ and $\alpha[C_i]$ is defined. Let

$$\widetilde{\eta}[u] = \alpha[\sigma] + \sum \pm\alpha[C_i],$$

$\pm$ is the sign of the end x_i of u. We now check this value is independent of choices.

Suppose we change a negative cap C to C', so the sphere σ changes to a new sphere σ'. Viewed in $\mathcal{L}_0 M$, the gluing $\sigma'' = C'\# - C$ is a sphere in M but also a sum of two 1-chains in $\mathcal{L}_0 M$, and under our identifications,

$$\alpha[\sigma''] = \alpha[C'] - \alpha[C].$$

In $\pi_2(M)$, the equality $\sigma'' + \sigma = \sigma'$ holds, so evaluating α we deduce

$$\alpha[C'] - \alpha[C] + \alpha[\sigma] = \alpha[\sigma'].$$

So $\alpha[\sigma] - \alpha[C]$ does not change when we replace the cap C with C'. Similarly for positive caps C. Thus $\widetilde{\eta}[u]$ only depends on u, not on the caps C_i.

Suppose we change the representative α by an exact form df, $f : \mathcal{L}_0 M \to \mathbb{Z}$. Then $SC^*(H)_\alpha \cong SC^*(H)_{\alpha+df}$ via the change of basis isomorphism

$$x \mapsto x' = t^{f(m)-f(x)}x.$$

We now check that this isomorphism respects the TQFT structure. Since $df[\sigma]=0$, $df[u] = \sum \pm df[C_i] = \sum \pm(f(x_i) - f(m)) = (p-q)f(m) + \sum \pm f(x_i)$. Hence, if $\psi(\otimes_b y_b) = \lambda\cdot\otimes_a x_a + \cdots$ computed in $SC^*(M)_\alpha$, then computed in $SC^*(M)_{\alpha+df}$:

$$\begin{array}{rcl} \psi(\otimes_b y'_b) & = & t^{qf(m)-\sum f(y_b)}\psi(\otimes_b y_b) \\ & = & t^{qf(m)-\sum f(y_b)}(t^{(p-q)f(m)-\sum f(x_a)+\sum f(y_b)}\lambda\cdot\otimes_a x_a + \cdots) \\ & = & \lambda\cdot\otimes_a x'_a + \cdots, \end{array}$$

as required. The construction of the TQFT for $SH^*_0(M)_{\widetilde{\eta}}$ is now routine. $\square$

Remark 21. *We used integral singular 1-cocycles α, but we can also use smooth real 1-cocycles in $\tau H^2(\widetilde{M};\mathbb{R}) \subset H^1(\mathcal{L}_0 M;\mathbb{R})$ via $H^1(\mathcal{L}\widetilde{M};\mathbb{R}) \cong \mathrm{Hom}(\pi_2 M,\mathbb{R})$.*

6.7. **Twisting by 1-forms from the base.** Let $u : S \to \overline{M}$ be an isolated Floer solution. For cylinders S we already defined $ev^*\mu[u]$ in 6.1. For a cap S, $ev^*\mu[u]$ cannot be defined consistently (it would not glue correctly). For other S, let S' be an associated graph (15.19). S' can be embedded into S so that the ends of S' ("exterior vertices") converge to $t=0\in S^1$ in the asymptotic circles in S. Associate to each edge e_i a closed 1-form μ_i on N, subject to the *sum-zero condition*: at any internal vertex v of the graph the signed sum of the forms vanishes,

$$\sum_{\substack{\text{edges } e_i \\ \text{coming into } v}} \mu_i - \sum_{\substack{\text{edges } e_i \\ \text{leaving } v}} \mu_i = 0.$$

Example. P' (Figure 12), assign 2μ to the incoming edge, μ to each outgoing edge.

Let u be a Floer solution $u : S \to \overline{M}$ or a Morse solution $u : S' \to \overline{M}$. Define

$$\mu[u] = \sum_{e_i} \mu_i[u|_{e_i}].$$

Lemma 22. *$\mu[u]$ is invariant under homotopying S' (subject to sum-zero).* □

Pick any $S' \subset S$ and μ_i as above. Counting Floer solutions with weight $t^{\mu[u]}$ defines

$$\psi_S : \bigotimes_{b=1}^{q} SH^*(M)_{ev^*\mu_b} \to \bigotimes_{a=1}^{p} SH^*(M)_{ev^*\mu_a} \qquad (p \geq 1, q \geq 0, (p,q) \neq (1,0)).$$

Lemma 23. *If the μ_a, μ_b satisfy the steady-state condition $\sum ev^*\mu_a = \sum ev^*\mu_b$, then it is possible to choose an S' satisfying the sum-zero condition.* □

By Lemma 22, ψ_S does not depend on the choice of graph S' associated to S. Gluing surfaces requires the additional condition that the forms μ_i agree on the gluing ends. Under this assumption, Theorem 91 follows immediately for these ψ_S.

Since $SH^*(M)_{ev^*(c\mu)} \cong SH^*(M)_{ev^*\mu}$ for any $c > 0$ (the local systems are isomorphic) one could use only multiples of $\mu \in H^1(M)$, which gives $SH^*(M)_{ev^*\mu}$ a TQFT structure without unit. More natural solutions are to define

$$NSH^*(M) = \bigoplus_{\mu \in H^1(M)} SH^*(M)_{ev^*\mu} \quad \text{and} \quad NSH^*(M)_\eta = \bigoplus_{\mu \in H^1(M)} SH^*(M)_{ev^*\mu + \tau\eta}.$$

with $c^* : \oplus_\mu NH^*(M)_\mu \to NSH^*(M)$ and $c^* : \oplus_\mu NH^*(M)_\mu \to NSH^*(M)_\eta$, where $NH^*(M)_\mu = H^*(M)_{ev^*\mu}$ is the Novikov cohomology of M. The products are respectively $SH_{ev^*\mu_1} \otimes SH_{ev^*\mu_2} \to SH_{ev^*(\mu_1+\mu_2)}$ and $SH_{\mu,\eta} \otimes SH_{\mu',\eta} \to SH_{\mu+\mu',\eta}$.

Theorem 24. *$NSH^*(M)$ is a TQFT with unit $\psi_C(1) \in SH^0(M)_0 = SH^0(M) \otimes \Lambda$. Similarly $NSH^*(M)_\eta$ is a TQFT with unit $\psi_C(1) \in SH^0(M)_\eta$.*

Remark 25. *$\oplus_{\alpha \in H^1(\mathcal{L}_0 M)} SH^*_0(M)_\alpha = \oplus_{\widetilde{\eta}} NSH^*_0(M)_{\widetilde{\eta}}$ is a TQFT. The $\alpha = ev^*\mu + \tau\widetilde{\eta}$ yield all classes in $H^1(\mathcal{L}_0 M)$: $\pi_1(\mathcal{L}_0 M) \cong \pi_1(M) \ltimes \pi_2(M)$ so dually $H^1(\mathcal{L}_0 M) \cong ev^* H^1(M) \oplus \mathrm{Hom}(\pi_2(M), \mathbb{Z})$. Also $c^* : \oplus_\alpha H^*(M)_\alpha \to \oplus_\alpha SH^*_0(M)_\alpha$.*

6.8. **Twisted wrapped theory.** Let $\alpha \in H^1(\Omega(M,L)) \cong H^1(\Omega(\overline{M},\overline{L}))$ (3.3). As in 6.1, α defines a local system of coefficients $\underline{\Omega}_\alpha$ on $\Omega(\overline{M},\overline{L})$. Define $CW^*(L;H)$ over the Novikov ring Λ and insert weights $t^{\alpha[u]}$ in the definitions of the differential and the continuation maps. This yields Λ-modules $HW^*(L;H;\underline{\Omega}_\alpha)$ and their direct limit $HW^*(L)_\alpha = HW^*(L;\underline{\Omega}_\alpha)$. These come with a canonical map

$$c^* : H^*(L; c^*\underline{\Omega}_\alpha) \to HW^*(L; \underline{\Omega}_\alpha),$$

where $c : \Omega(\overline{L},\overline{L}) \to \Omega(\overline{M},\overline{L})$ is the inclusion of paths lying in $\overline{L}$.

Let $\eta \in H^2(\overline{M},\overline{L};\mathbb{R}) \cong H^2(M,L;\mathbb{R}) \cong H^2(M/L;\mathbb{R})$. One can represent η by a closed de Rham two-form which is supported away from $\overline{L}$ (see [21, Sec.11.1]). Evaluating $\int_C \sigma^*\eta$ on smooth one-chains $\sigma : C \to \overline{M}$ defines a transgresssion cycle $\tau\eta \in H^1(\Omega(\overline{M},\overline{L}))$, which can also be obtained via: (using real coefficients)

$$\tau : H^2(M/L) \overset{ev^*}{\to} H^2(\Omega(M/L)\times S^1) \overset{\text{project}}{\longrightarrow} H^1(\Omega(M/L)) \cong \mathrm{Hom}(\pi_2(M/L),\mathbb{R}).$$

This is an isomorphism when $\pi_1(M/L) = 1$ (e.g. if $\pi_1(M) = 1$). For simply connected L, M, τ is the Hurewicz isomorphism $H^2(M,L;\mathbb{R}) \cong \mathrm{Hom}(\pi_2(M,L),\mathbb{R})$.

For a wrapped solution $u : S \to \overline{M}$ define $\tau\eta[u] = \int_S u^*\eta$. These weights are locally constant on the moduli spaces of wrapped solutions since $\tau\eta[u]$ remains constant if we homotope u relative to the ends while keeping $u(\partial S) \subset \overline{L}$. Therefore, denoting $\overline{\eta} \in H^2(M;\mathbb{R})$ the image of η via $H^2(M,L;\mathbb{R}) \to H^2(M;\mathbb{R})$,

Theorem 26. *$HW^*(L;\underline{\Omega}_{\tau\eta})$ is a module over $SH^*(M;\underline{\Lambda}_{\tau\overline{\eta}})$ and has a TQFT structure and $c^* : H^*(L)\otimes\Lambda \to HW^*(L;\underline{\Omega}_{\tau\eta})$ respects the TQFT.*

7. Viterbo functoriality

7.1. **No escape lemma.** To prove compactness results, the following lemma will be useful (it is analogous to Abouzaid-Seidel [3, Lemma 7.2], adapted to our setup).

Let $(V, d\theta)$ be a symplectic manifold with boundary ∂V of negative contact type: the Liouville field Z, defined by $i_Z d\theta = \theta$, points strictly inwards. Near ∂V define the coordinate $R = e^r$ parametrizing the time r flow of Z starting from $\partial V = \{R = 1\}$ (1.1). Assume J is of contact type on ∂V: $J^*\theta = dR$. Suppose $H : V \to [0,\infty)$ is linear near ∂V, say $H = mR$. Define X by $i_X d\theta = -dH$. Let S be a compact Riemann surface with boundary with a 1-form β such that $d\beta \leq 0$.

Lemma 27. *Any solution $u : S \to V$ of $(du - X\otimes\beta)^{0,1} = 0$, with $u(\partial S) \subset \partial V$, will map entirely into ∂V and solve $du = X\otimes\beta$.*

Proof. We use a trick we learnt from Mohammed Abouzaid: we show that $E(u) \leq 0$. Therefore $E(u) = 0$, so $du = X\otimes\beta$, so du lands in the span of $X = h'\mathcal{R} \subset T\partial V$, thus $u(S) \subset \partial V$ as required. On ∂V, $X = h'(R)\,\mathcal{R}$, so $\theta(X) = mR = H$, thus

$$\begin{array}{rcll}
E(u) & = & \int_S u^*d\theta - u^*dH\wedge\beta & (\text{Section 15.3 using } \omega = d\theta) \\
& \leq & \int_S du^*\theta - d(u^*H\beta) & (d\beta \leq 0 \text{ and } H \geq 0) \\
& = & \int_{\partial S} u^*\theta - u^*H\beta & (\text{Stokes's Theorem}) \\
& = & \int_{\partial S} u^*\theta - \theta(X)\beta & (H = \theta(X) \text{ on } u(\partial S) \subset \partial V) \\
& = & \int_{\partial S} \theta(du - X\otimes\beta) & \\
& = & \int_{\partial S} -\theta J(du - X\otimes\beta)j & (\text{holomorphicity of } du - X\otimes\beta) \\
& = & \int_{\partial S} -dR(du - X\otimes\beta)j & (J \text{ is contact on } u(\partial S) \subset \partial V) \\
& = & \int_{\partial S} -dR(du)j & (dR \text{ vanishes on } X = m\mathcal{R})
\end{array}$$

Let $\hat{n}$ = outward normal direction along $\partial S \subset S$. Then $(\hat{n}, j\hat{n})$ is an oriented frame, so ∂S is oriented by $j\hat{n}$. Now $dR(du)j(j\hat{n}) = d(R\circ u)\cdot(-\hat{n}) \geq 0$ since in the inward direction $-\hat{n}$, $R\circ u$ can only increase since ∂V minimizes R. So $E(u) \leq 0$. ☐

7.2. **Liouville subdomains.** A Liouville subdomain $i : (W^{2n},\theta_W) \hookrightarrow (M^{2n},\theta)$ is the embedding of a Liouville domain (W,θ_W) such that for some $\rho \in \mathbb{R}$,

$$i^*\theta = e^\rho\theta_W + \text{exact form}.$$

Example 28. *Suppose* $L^n \subset DT^*N$ *is an exact Lagrangian submanifold* ($\theta|_L =$ *exact*)*. This induces a Weinstein embedding* $i : DT^*L \hookrightarrow DT^*N$ *of a small disc cotangent bundle of* L *onto a neighbourhood of* L*, and* i *is a Liouville subdomain.*

Lemma 29. *For the purposes of symplectic cohomology, one can always assume a Liouville embedding is an inclusion* $W \subset M$*, using the same* θ, Z *for* W *as for* M*.*

Proof. We can always assume $\rho = 1$ by redefining θ_W to be $e^{\rho}\theta_W$. This does not affect the symplectic cohomology theory for $\overline{W}$ (it rescales all Hamiltonians by e^{ρ}).

Identify W and $i(W)$. We are given $\theta_W = \theta - df$, with $f : W \to \mathbb{R}$. Extend f smoothly to $\overline{M}$, so that it vanishes outside of a compact neighbourhood of W. Then extend θ_W, Z_W to all of $\overline{M}$ by $\theta_W = \theta - df$ and $\omega(Z_W, \cdot) = \theta_W$.

Since $d\theta_W = d\theta$ everywhere and $\theta_W = \theta$ on the collar of $\overline{M}$, the Hamiltonian vector fields agree and we can use the same almost complex structure. So the Floer theory is the same: identify $SH^*(M, \theta_W) = SH^*(M, \theta)$, and replace θ by θ_W. $\square$

Remark 30. *Any Liouville embedding* $i : (W, \theta_W) \hookrightarrow (M, \theta_M)$ *can now be extended to an embedding* $i : (\overline{W}, \theta_W) \hookrightarrow (\overline{M}, \theta_M)$ *by defining* i *on the collar to be the Liouville flow of* Z_W *in* $\overline{M}$ (*so* $[1, \infty) \times \partial W \ni (R = e^r, y) \mapsto \varphi^r_{Z_W}(y) \in \overline{M}$)*. Since* Z_W *is outward pointing along* ∂W*, this flow will not reintersect* W*.*

By Lemma 29, we always assume $W \subset M$. Let $C_W = [1, 1+\varepsilon) \times \partial W$ be a collar of $\partial W \subset \overline{M}$. Choose J on $\overline{M}$ of contact type near $R = e^r = 1 + \varepsilon$. Choose $H : \overline{M} \to \mathbb{R}$ with $H = mR$ near $R = 1 + \varepsilon$ and $H \geq 0$ on $\overline{M} \setminus (W \cup C_W)$.

Corollary 31. *Any Floer solution with asymptotics in* $W \cup C_W$ *cannot escape* W*.*

Proof. Choose $V = \overline{M} \setminus (W \cup [1, R) \times \partial W)$, with R close to $1+\varepsilon$ chosen generically so that $\partial V = \{R\} \times \partial W$ intersects u transversely. Apply Lemma 27 to the restriction of u to the preimage of each component of $u(S) \cap V$. Hence $u(S) \subset W \cup C_W$. $\square$

7.3. **Viterbo Functoriality.** For Liouville subdomains $W \subset M$, Viterbo [36] constructed a restriction map $SH^*(M) \to SH^*(W)$ and McLean [22] observed that it is a ring homomorphism. We now prove a stronger statement:

Theorem 32. *Let* $i : (W, \theta_W) \hookrightarrow (M, \theta)$ *be a Liouville subdomain. Then there exists a TQFT map called* restriction map*,* $SH^*(i) : SH^*(M) \to SH^*(W)$*, which fits into a commutative diagram which respects TQFT structures:*

$$\begin{tikzcd}
SH^*(W) & SH^*(M) \arrow[l, "SH^*(i)"'] & & SH^*(W)_{i^*\eta} & SH^*(M)_{\eta} \arrow[l, "SH^*(i)_{\eta}"'] \\
H^*(W) \arrow[u, "c^*"'] & H^*(M) \arrow[l, "i^*"'] \arrow[u, "c^*"'] & & H^*(W) \otimes \Lambda \arrow[u, "c^*"'] & H^*(M) \otimes \Lambda \arrow[l, "i^*"'] \arrow[u, "c^*"']
\end{tikzcd}$$

In particular, all maps are unital ring homomorphisms.

Proof. We may assume $W \subset \overline{M}$ and $\theta_W = \theta|_W$ by Lemma 29. Pick a (smoothing of the) diagonal-step shaped Hamiltonian H in Figure 2. More precisely, the Liouville flow parametrizes a tubular neighbourhood $(-\infty, 1+\varepsilon) \times \partial W$ of $\partial W \subset M$. Using (in general different) R-coordinates on the two collars, choose H linear of slope b on the collar of $\overline{M}$ and linear of slope a (with $a \gg b$) on $(c, 1+\varepsilon) \times \partial W$. Interpolate H smoothly so that it is constant on $M \setminus W$ and on $W \setminus (c, 1] \times \partial W$ except near their boundaries. Here a, b are generic (not Reeb periods). It was carefully proved in [27] that there is a $\delta > 0$ such that for $c \ll a^{-1}$ the 1-periodic orbits in $W \cup (1, 1+\varepsilon) \times \partial W$ have actions $A_H < \delta$, and all orbits outside that region have actions $A_H > \delta$.



FIGURE 2. A diagonal-step shaped Hamiltonian with slopes a, b. The dashed line is the action $A(R) = -Rh'(R) + h(R)$.

Since the action decreases along Floer trajectories, we have a commutative diagram of action-restriction chain maps,

$$\begin{tikzcd}
CF^*(M,H';A_{H'}<\delta)_\eta & CF^*(M,H')_\eta \arrow[l] \\
CF^*(M,H;A_H<\delta)_\eta \arrow[u] & CF^*(M,H)_\eta \arrow[l] \arrow[u]
\end{tikzcd}$$

where on the left the chain complex is generated only by those orbits which have action less than δ. Here H' is another diagonal-step shaped Hamiltonian with slopes $a' > a$, $b' > b$ such that $H' > H$, and the vertical maps are the (twisted) continuation maps corresponding to a monotone homotopy from H' to H.

We claim that the commutative diagram of chain maps below respects the TQFT,

$$\begin{tikzcd}
CF^*(W,i^*H')_{i^*\eta} & CF^*(M,H';A_{H'}<\delta)_\eta \arrow[l, "i^{-1}"'] \\
CF^*(W,i^*H)_{i^*\eta} \arrow[u] & CF^*(M,H;A_H<\delta)_\eta \arrow[l, "i^{-1}"'] \arrow[u]
\end{tikzcd}$$

For this, we need Corollary 31 three times. Firstly, Floer trajectories with ends in W do not escape $R \leq 1+\varepsilon$, hence the i^{-1} maps are well-defined identifications and they are chain maps. Secondly, Floer continuation solutions do not escape, so the i^{-1} commute with the vertical continuation maps. Thirdly, the TQFT operations on the above groups involve counting Floer solutions with ends in W, and these cannot escape $R \leq 1+\varepsilon$, so i^{-1} respects the TQFT structure.

So we obtain the following commutative diagrams respecting TQFT structures

$$\begin{tikzcd}
SH^*(W,i^*H')_{i^*\eta} & SH^*(M,H')_\eta \arrow[l] \\
SH^*(W,i^*H)_{i^*\eta} \arrow[u] & SH^*(M,H)_\eta \arrow[l] \arrow[u]
\end{tikzcd}$$

Take $H = H^\delta$ as in 4, identify $SH^*(M,H^\delta)_\eta \cong H^*(M)\otimes\Lambda$, and take the direct limit over the H' to obtain the claim using Theorems 14, 17. □

Remark 33.

(1) *The restriction map* $SH^*(i) : SH^*(M)_\eta \to SH^*(W)_{\eta|W}$ *is independent of the choice of embedding* $i : W \hookrightarrow \overline{M}$ *satisfying* $i^*\theta_M - e^{constant}\theta_W = exact$*.*

(2) *Functoriality: given nested Liouville subdomains* $W' \hookrightarrow W \hookrightarrow \overline{M}$*, then* $SH^*(M)_\eta \to SH^*(W)_{i^*\eta} \to SH^*(W')_{i'^*\eta}$ *equals* $SH^*(M)_\eta \to SH^*(W')_{i'^*\eta}$*.*

(3) *For* $i = id|_M$ *or any inclusion* $i : M \cup [1,R_0]\times\partial M \hookrightarrow \overline{M}$*,* $SH^*(i) = id$*.*

Sketch proof. (1) Consider a family i_λ, $0 \leq \lambda \leq 1$, of such embeddings. By Lemma 29, this reduces to considering a fixed $W \subset \overline{M}$ with $\theta = \theta_\lambda$ varying only in a compact neighbourhood of W. Then one checks that the restriction maps $SH^*(i_0)$ and $SH^*(i_1)$ are chain homotopic, for example by using the 1-parameter family of moduli spaces of Floer trajectories for θ_λ and suitable step-shaped H_λ.

(2) Using an appropriate step-shaped Hamiltonian with two steps instead of one, one can separate the action values of the generators in W', $W \setminus W'$ and $\overline{M} \setminus W$.

(3) For $i = \mathrm{id}$ one checks directly that the map induced by the step-shaped Hamiltonian is the continuation map $SH^*(H^b) \to SH^*(H^a)$ in the notation 2.3, so the direct limit as $a \gg b \to \infty$ is the identity. For inclusions, use the Liouville flow to push $M \cup [1, R_0] \times \partial M$ onto M and use (1) to reduce to the $i = \mathrm{id}$ case. □

7.4. **Wrapped Viterbo restriction.** Let $W \hookrightarrow M$ be a Liouville subdomain, and L an exact Lagrangian with transverse Legendrian intersections with $\partial W, \partial M$ (by Lemma 29, we can assume $W \subset M$, $\theta_W = \theta|_W$). So explicitly: $\theta|_L = df$ and $df = 0$ on $L \cap (\partial W \cup \partial M)$. To define a restriction map, we need [3, Ex.4.2] an additional assumption: *one can extend $f : L \to \mathbb{R}$ to $f : \overline{M} \to \mathbb{R}$ such that f is locally constant on $\partial W \cup \partial M$.* Then the analogue of Theorem 32 holds:

Theorem 34. *There is a restriction map $HW^*(L \subset M) \to HW^*(L \cap W \subset W)$, commuting with c^*, and compatible with the TQFT and the SH^*-module structure. More generally there is a twisted restriction $HW^*(L; \underline{\Omega}_\alpha) \to HW^*(L \cap W; \underline{\Omega}_{\alpha|W})$, which for $\alpha = \tau\eta$ respects the structures of Theorem 26.*

Proof. We need $f|_L = 0$ near $\partial W \cup \partial M$. This can be achieved after deforming L by a Hamiltonian isotopy of M relative to $\partial W \cup \partial M$ ([3, Lemma 4.1]). It ensures L has the form (interval) × $(L \cap \partial W)$ near ∂W and similarly near ∂M.

Now proceed as in 7.3. Suppose for now that we can separate the action values as in the proof of Theorem 32. Then the proof is analogous using $CW^*(L; H)$, $CW^*(L \cap W; H)$, provided Corollary 31 holds: this we now discuss.

We need the analogue of Lemma 27 for: ∂V a generic level set $R_W = 1 + \varepsilon$ near ∂W; V the unbounded component of $\overline{M}$ outside of that level set; and S the compact Riemann surface with boundary obtained as the preimage of a component of $V \cap \mathrm{image}(u)$ for a wrapped solution u whose ends are Hamiltonian chords lying inside W. From now on, u denotes the restriction $u|_S$.

As in Lemma 27, we need $E(u) \leq 0$. The boundary ∂S contains not just $S \cap \partial V$ (which are now circles and line segments) but also boundary pieces lying in $\overline{L} \cap V$. So Stokes applied to $E(u) \leq \int_{\partial S} u^*\theta - u^*H\beta$ yields new contributions coming from curves $\gamma \subset \partial S$ with $u(\gamma) \subset \overline{L}$. The $u^*H\beta$ gives no new contributions since $\beta|_\gamma = 0$, but the $u^*\theta|_{\overline{L}} = u^*df$ term produces new summands of the form $f(z) - f(z')$ for $z, z' \in \partial V \cap \overline{L}$. For small ε, $\overline{L} \cap \partial V$ is so close to $\overline{L} \cap \partial W$ that $f|_{\overline{L}} = 0$, and so $f(z) = f(z') = 0$. Thus $E(u) \leq 0$ still holds.

We now show how to separate the action values. The action on generators is $-Rh'(R) + h(R)$ provided $f = 0$ (3.3). Thus if generators lie where $f = 0$ then the action calculations for SH^* hold also for HW^*, and we are done. It remains to ensure $f = 0$ near $R_W = c$ (where W generators lie).

Let $\varphi = \varphi_Z^{\log c}$ = Liouville flow for time $\log c < 0$. Push $L \cap W$ into $\varphi(L \cap W)$. Since L has the form (interval) × $(L \cap \partial W)$ near ∂W, the flow rescales the interval by c, so we can smoothly join $\varphi(L \cap W)$ to $\overline{L} \setminus W$ by some (interval) × $(L \cap \partial W)$.

Call P the resulting Lagrangian. For (P,H,W,M,θ) actions can be separated so

$$HW^*(P;H) \overset{\text{restrict}}{\longrightarrow} HW^*(P;H,A_H<\delta) \cong HW^*(W\cap P;H|_W).$$

We claim that this naturally induces a restriction $HW^*(L;H^b)\to HW^*(W\cap L;H^a)$ for some H^b,H^a with slopes b,a at infinity on $\overline{M},\overline{W}$. This claim implies the Theorem after taking direct limits $a\gg b\to\infty$. Since $\overline{L},P$ are exact and isotopic, there is a compactly supported Hamiltonian isotopy of $\overline{M}$ mapping $\overline{L}$ onto P. This induces an isomorphism $HW^*(P;H)\cong HW^*(L;H^b)$ by pull-back. Next, φ induces

$$\varphi^*: HW^*(W\cap P;H|_W;\theta)\cong HW^*(W\cap L;\varphi^*(H|_W);c\,\theta).$$

where $\varphi^*(H|_W)=H|_W\circ\varphi$ has slope ac at infinity on $\overline{W}$. Rescaling $c\,\theta$ and $\varphi^*(H|_W)$ by $1/c$ preserves the Hamiltonian vector field and thus the whole Floer theory:

$$HW^*(W\cap L;\varphi^*(H|_W);c\,\theta)\cong HW^*(W\cap L;H^a;\theta),$$

where $H^a=\varphi^*(H|_W)/c$ has slope a at infinity, as required. $\square$

8. Vanishing Criteria

Theorem 35. *1) $SH^*(M)_\eta=0$ if and only if the unit $e=0$.*
2) For Liouville $i:(W,\theta_W)\hookrightarrow(M,\theta)$, if $SH^*(M)_\eta=0$ then $SH^*(W)_{i^*\eta}=0$.

Proof. 1) If $e_M=0$ then for $y\in SH^*(M)_\eta$, $y=e_M\cdot y=0\cdot y=0$, so $SH^*(M)_\eta=0$.
2) Suppose $SH^*(M)_\eta=0$, so $e_M=0$. The restriction $\varphi:SH^*(M)_\eta\to SH^*(W)_{i^*\eta}$ is a TQFT map, so $e_W=\varphi(e_M)=\varphi(0)=0$. $\square$

Theorem 36. *$SH^*(M)=0$ iff $e_H=0$ for some H iff $c^*:H^*(M)\to SH^*(M)$ vanishes iff $c^*:H^*(M)\to SH^*(H)$ vanishes for some H. Similarly for $SH^*(M)_\eta$.*

Proof. $e=\varinjlim e_H=0$ occurs by definition iff some $e_H=0$. If $e_H=0$, then using $\psi_P:H^*(M)^{\otimes 2}\to SH^*(H)$ it follows that $0=\psi_P(e_H,y)=c^*y\in SH^*(H)$ for all $y\in H^*(M)$. So $e_H=0$ implies $c^*:H^*(M)\to SH^*(H)$ vanishes, and the converse follows because $e_H=c^*(1)$. Note that $c^*:H^*(M)\to SH^*(H)\to SH^*(M)$ factors, and the rank of the image in $SH^*(H)$ is finite and decreasing as the slope of H at infinity increases. So the rank eventually stabilizes. This proves the last 'iff'. $\square$

Dualizing the above proof yields Theorem 37. Using that c^*,c_* are dual to each other yields Theorem 38. Arguments analogous to the above prove Theorem 39.

Theorem 37. *$SH_*(M)=0$ iff the counit $\psi^C:SH_*(M)\to\mathbb{K}$ vanishes iff $c_*:SH_*(M)\to H_*(M)$ vanishes iff they vanish for some H. Similarly for $SH_*(M)_\eta$.*

Theorem 38. *$SH^*(M)=0$ iff $SH_*(M)=0$.*

Theorem 39. *$HW^*(L)=0$ iff the unit $\mathcal{W}_C(1)=0$ iff $c^*:H^*(L)\to HW^*(L)$ vanishes iff they vanish for some H. Similarly for $HW^*(L)_\eta$.*

Theorem 40. *$SH^*(M)=0\Rightarrow HW^*(L)=0$, and $SH^*(M)_{\overline{\eta}}=0\Rightarrow HW^*(L)_\eta=0$.*

Proof. $e=0$ in SH^*, so $y=e\cdot y=0\cdot y=0$ for $y\in HW^*$ (Theorems 101, 26). $\square$

9. THE SH^*_+ AND HW^*_+ GROUPS.

In the following proof, we define the groups $SH^*_+(M)$ (twisted by $\alpha \in H^1(\mathcal{L}M)$, when twistings are present). The untwisted construction is well-known [36, 7].

Theorem 41. *There is a long exact sequence induced by action-restriction maps*

$$\cdots \to H^*(M)_\alpha \xrightarrow{c^*} SH^*(M)_\alpha \to SH^*_+(M)_\alpha \to H^{*+1}(M)_\alpha \to \cdots$$

Proof. The orbits inside M are critical points of H with action close to zero, say $A_H \in (-\delta, \delta)$. On the collar, pick H convex and growing slowly until it becomes linear at infinity. The first orbit on the collar appears when H has slope equal to the smallest Reeb period. By making the growth of H slow enough, we can make $A_H = -Rh'(R) + h(R)$ arbitrarily negative on orbits on the collar, so $A_H < -\delta$.

Write $SC^*(A,B)_\alpha$ for the chain complex generated only by the orbits with action in the interval (A,B) (for generic $A, B \in \mathbb{R} \cup \{\pm\infty\}$, so not Reeb periods).

Action decreases along Floer trajectories, so increasing A gives a subcomplex $SC^*(A',B)_\alpha \hookrightarrow SC^*(A,B)_\alpha$ for $A < A' < B$ and decreasing B gives quotient maps $SC^*(A,B)_\alpha \to SC^*(A,B')_\alpha$ for $A < B' < B$. So $A < A' = B' < B$ gives the exact

$$0 \to SC^*(A',B)_\alpha \hookrightarrow SC^*(A,B)_\alpha \to SC^*(A,A')_\alpha \to 0.$$

This induces a long exact sequence, which for $-\infty < -\delta < \delta$ is

$$\cdots \to SH^*(-\delta,\delta)_\alpha \xrightarrow{c^*} SH^*(-\infty,\delta)_\alpha \to SH^*(-\infty,-\delta)_\alpha \to SH^{*+1}(-\delta,\delta)_\alpha \to \cdots$$

where c^* is the inclusion of the Morse subcomplex (Section 4).

Definition: *$SH^*_+(H)_\alpha = SH^*(-\infty,-\delta)_\alpha$, and $SH^*_+(M)_\alpha$ is their direct limit.* $SH^*(-\delta,\delta)_\alpha \cong H^*(M)_\alpha$, $SH^*(-\infty,\delta)_\alpha = SH^*(H)_\alpha$. Now take direct limits. □

By [27, Thm 3] the isomorphism $SH^*(T^*N)_\alpha \cong H_{n-*}(\mathcal{L}N)_\alpha$ respects certain action filtrations, which for the action bounds $(-\infty,-\delta)$ above yields

$$SH^*_+(T^*N)_\alpha \cong H_{n-*}(\mathcal{L}N, N)_\alpha.$$

Corollary 42. *For $\overline{M} = T^*N$ the long exact sequence is that of the pair $(\mathcal{L}N, N)$:*

$$\begin{tikzcd}
\cdots \arrow[r] & H^*(T^*N)_\alpha \arrow[r, "c^*"] \arrow[d, "\wr"] & SH^*(T^*N)_\alpha \arrow[r] \arrow[d, "\wr"] & SH^*_+(T^*N)_\alpha \arrow[r] \arrow[d, "\wr"] & \cdots \\
\cdots \arrow[r] & H_{n-*}(N)_\alpha \arrow[r, "c_{n-*}"] & H_{n-*}(\mathcal{L}N)_\alpha \arrow[r] & H_{n-*}(\mathcal{L}N, N)_\alpha \arrow[r] & \cdots
\end{tikzcd}$$

Theorem 43. *There is a long exact sequence for wrapped Floer cohomology,*

$$\cdots \to H^*(L)_\alpha \xrightarrow{c^*} HW^*(L)_\alpha \to HW^*_+(L)_\alpha \to H^{*+1}(L)_\alpha \to \cdots$$

Proof. The construction is analogous to Theorem 41. The additional term $f(x(1)) - f(x(0))$ in the action will not affect the $A_H < -\delta$ estimate on the collar, since we can make the other term $-Rh'(R) + h(R)$ arbitrarily small as in Theorem 41. □

10. ARNOL'D CHORD CONJECTURE

The Chord Conjecture states that a contact manifold containing a Legendrian submanifold has a Reeb chord with ends on the Legendrian. Our setup 3.1 involves the Legendrian $\partial L = L \cap \partial M$ inside the exact contact $\partial M \subset \overline{M}$.

Theorem 44. *If $HW^*(L) = 0$ then the Arnol'd chord conjecture holds for $\partial L \subset \partial M$ (existence of a Reeb chord). For a generic contact form α there are at least rank $H^*(L)$ chords. The same holds if $HW^*(L)_\eta = 0$ for $\eta \in H^2(M, L; \mathbb{R})$.*

Proof. Suppose there are no Reeb chords. Then we do not need to perturb α to avoid transversality issues caused by degenerate Reeb chords. By Lemma 8 and the maximum principle, $c^*: H^*(L) \cong HW^*(L; H^\delta) \cong HW^*(L)$, so $HW^*(L) \neq 0$.

Now suppose $HW^*(L) = 0$, α generic. By Theorem 43 if $HW^*(L) = 0$ then $HW^*_+(L) \cong H^{*+1}(L)$, so there are at least rank $H^*(L)$ distinct Hamiltonian chords on the collar (15.22.(3)), and the same for Reeb chords (Lemma 8). The twisted case is analogous, since $HW^*(L)_\eta = 0$ implies $HW^*_+(L)_\eta \cong H^{*+1}(L) \otimes \Lambda$. □

Remark 45. *The proof is similar to Viterbo's applications* [36] *to the Weinstein conjecture (existence of a closed Reeb orbit): if there are no closed Reeb orbits then $c^*: H^*(M) \to SH^*(M)$ is an isomorphism. So this conjecture holds if $SH^*(M) = 0$.*

In general rank $H^*(L) \geq \frac{1}{2}$rank $H^*(\partial L)$, using the long exact sequence for the pair $(L, \partial L)$ and observing that rank $H^*(L) =$ rank $H^*(L, \partial L)$ by Poincaré duality.

Example 46. *(Known to experts) Let M be a subcritical Stein manifold. Then Theorem 44 applies: $SH^*(M) = 0$ by Cieliebak* [10] *so $HW^*(L) = 0$ by Theorem 40.*

Let N be a simply connected closed manifold, and $L \subset DT^*N$ an exact Lagrangian with transverse Legendrian intersection $\partial L = L \cap ST^*N$.

Theorem 47. *If $H^2(T^*N) \to H^2(L)$ is not injective, then the chord conjecture holds and generically there are at least rank $H^*(L)$ Reeb chords.*

Proof. The non-injectivity implies the existence of an $\eta \in H^2(T^*N, L)$ with $\overline{\eta} \neq 0 \in H^2(T^*N)$. Combining Lemma 40 and the vanishing $SH^*(T^*N)_{\overline{\eta}} = 0$ (13.6) implies $HW^*(L)_\eta = 0$. Thus the claim follows by Theorem 44. □

Theorem 47 applies for example for L a conormal bundle to a submanifold $K \subset N$ such that $H^2(N) \to H^2(K)$ has kernel (Example 7).

Theorem 47 also holds after attaching subcritical handles to DT^*N, since Cieliebak [10] proved SH^* does not change for action reasons, so it also holds for twisted SH^*.

Theorem 48. *For any ALE space the chord conjecture holds for any ∂L and generically there are at least rank $H^*(L)$ Reeb chords.*

Proof. By [28], $SH^*(M)_{\overline{\eta}} = 0$ for generic $\overline{\eta} \in H^2(M)$. Now L is a 2-manifold with boundary so $H^2(L) = 0$, so $\overline{\eta}$ lifts to an $\eta \in H^2(M, L)$. Now see the proof of 47. □

11. Exact contact hypersurfaces

11.1. **Exact contact hypersurfaces.**

Definition 49. *An exact contact hypersurface is an embedding $i : (\Sigma^{2n-1}, \xi) \hookrightarrow (\overline{M}^{2n}, d\theta)$ of a closed contact manifold (Σ, ξ) such that for some 1-form $\alpha \in \Omega^1(\Sigma)$,*

$$\xi = \ker \alpha \quad \text{and} \quad \alpha - i^*\theta = \text{exact}.$$

Moreover, we will always assume that $i(\Sigma)$ separates M into a compact connected submanifold $W \subset \overline{M}$ and an unbounded component $\overline{M} \setminus W$.

Remark 50. *If $H_{2n-1}(M) = 0$, the separating condition is automatic: if $\overline{M} \setminus \Sigma$ is connected, a path γ in $\overline{M}$ cutting Σ once transversely has intersection $[\gamma] \cdot [\Sigma] \neq 0$ so $[\Sigma] \neq 0 \in H_{2n-1}(\overline{M})$. If $H_1(\Sigma) = 0$, then $\alpha - i^*\theta =$ exact is equivalent to $d\alpha = i^*\omega$.*

Example 51. *Sphere bundles* $ST^*N \hookrightarrow T^*N$*, boundaries* $ST^*L \hookrightarrow M$ *of Liouville domains of Example 28* ($n \geq 2$ *for connectedness*)*. The separating condition for* $\Sigma \hookrightarrow T^*N$ *is automatic if* $dim(N) = n \geq 2$*, since* $H_{2n-1}(T^*N) \cong H_{2n-1}(N) = 0$*.*

Lemma 52. *For the purposes of symplectic cohomology, we can always assume that an exact contact hypersurface* Σ *bounds a Liouville subdomain* W *of* M*.*

Proof. Assume $i(\Sigma) \subset M$ by redefining M to be $M \cup [1, R_0] \times \partial M$ (Remark 33). Identify $i(\Sigma) = \Sigma$. Use a bump function to extend α to a neighbourhood of $\Sigma \subset \overline{M}$. Near Σ, $\alpha - \theta = df$ for a function f supported near Σ. Then replace θ by $\theta + df$: $SH^*(M)$ has not changed (compare Lemma 29). Thus we can assume $W \subset M$ with $\alpha = \theta|_\Sigma$ a contact form. Define Z by $\omega(Z, \cdot) = \theta$, so $\omega(Z, \cdot)|_\Sigma = \alpha$. Thus, $\omega^n(Z, \ldots)|_\Sigma = \alpha \wedge (d\alpha)^{n-1} \neq 0$ pointwise (contact condition) and since the flow of Z expands volumes, Z points strictly outward along Σ. So $W \subset M$ is Liouville. ☐

11.2. Stretching-of-the-neck argument.

We recall this argument of Bourgeois-Oancea [7, Sec.5.2]. Consider an isolated Floer trajectory in $\overline{M}$ joining orbits lying in the collar. Consider what happens to this Floer trajectory as you stretch a neighbourhood of ∂M (more precisely: you insert a collar $[R_0, 1] \times \partial M$ in between M and $[1, \infty) \times \partial M$, and you rescale ω, H on M by $1/R_0$ so that the data glues correctly). By a compactness argument, this 1-family of Floer trajectories will converge in the limit to a Floer cylinder with punctures

$$(\mathbb{R} \times S^1) \setminus \{\text{punctures}\} \to \partial M \times \mathbb{R}$$

and the map rescaled near the punctures converges to Reeb orbits, each of which is "capped off" by an isolated holomorphic plane $\mathbb{C} \to \overline{M}$ (converging to the Reeb orbit at infinity). The proof that the only holomorphic curves capping off the Reeb orbits in $\overline{M}$ are planes (and not, for example, holomorphic cylinders) is a consequence of a subtle action argument [7, Sec.5.2, Proof of Prop.5, Step 1] which shows that the limit curve must have a *connected* component containing both Hamiltonian orbits (the fixed ends of the Floer trajectory). Combining this with a dimension count and the fact that cylinders have genus 0 proves the above are the only limits.

Remark 53 (Transversality)**.** *In general, this argument would require knowing transversality for symplectic field theory* [7, Remark 9, sec.3.1]*. In our applications this is not necessary, as explained to us by Oancea: we will always assume that the virtual dimension* $(2n - 3) - |x|$ *of the moduli space of holomorphic planes* $\mathbb{C} \to \overline{M}$ *converging to a Reeb orbit* x *at infinity is* ≥ 1*, therefore the above punctured Floer cylinders do not have punctures. Indeed, if they had punctures, then since the virtual dimension of the planes is* ≥ 1*, the virtual dimension of the component containing the two Hamiltonian orbits (modulo the* $\mathbb{R}$*-translation action) would be negative. But for that component, transversality is guaranteed by a time-dependent perturbation of the almost complex structure just like for non-punctured Floer cylinders. Thus that component does not exist, which is a contradiction.*

11.3. Independence of the filling.

Lemma 54 (Bourgeois-Oancea [7], Cieliebak-Frauenfelder-Oancea's Thm 1.14 [12])**.** *Let* W *be a Liouville domain with* $c_1 = 0$ *or* $c_1(\pi_2(W)) = 0$ *such that all closed Reeb orbits* x *in* ∂W *which are contractible in* W *have* $|x| < 2n - 3$*. Then for small* $C > 0$*, all isolated Floer trajectories in* $\overline{W}$ *connecting orbits in the collar lie in* $R \geq C$*.*

For a disc cotangent bundle $W = DT^*L$ *these conditions are satisfied if* $\dim L \geq 4$*.*

Proof. If no such C existed, 11.2 implies the existence of an isolated punctured Floer cylinder capped off by at least one holomorphic plane in $\overline{W}$. But $(2n-3)-|x|\geq 1$, so Remark 53 implies that the cylinder cannot have punctures, contradiction.

For $W^{2n}=DT^*L$, $c_1(W)=0$. Our grading convention is $SH^*(T^*L)\cong H_{n-*}(\mathcal{L}L)$, so $|x|=n-\mathrm{index}(\gamma_x)$ where γ_x is the closed geodesic (Reeb orbit) corresponding to x. So $(2n-3)-|x|=\mathrm{index}(\gamma_x)+\dim L-3$. The claim follows since $\mathrm{index}(\gamma_x)\geq 0$. $\square$

Corollary 55. *1) The* boundary symplectic cohomology

$$BSH^*(\partial W)_{\eta|\partial W}=SH^*_+(W)_\eta \qquad (\textit{Section 9})$$

is independent of the choice of filling W of ∂W which satisfies the conditions of Lemma 54. In particular, it only depends on the restriction of η to $H^2(\partial W)$.

2) Let $i:\Sigma\hookrightarrow\overline{M}$ be an exact contact hypersurface with $c_1(M)=0$ and all closed Reeb orbits x in Σ having $|x|<2n-3$ in $\overline{M}$. Then $BSH^*(\Sigma)_{\eta|\Sigma}\cong SH^*_+(W)_\eta$ for the Liouville subdomain W bounding $i(\Sigma)$ in $\overline{M}$ (Lemma 52).

Proof. 1) We can choose the same Hamiltonian and almost complex structure on $R\geq C$ for two such fillings, so the Floer equations are identical. Twistings are determined by η restricted to $R\geq C$, but this region is homotopy equivalent to ∂W.

2) $c_1(M)=0$, so $c_1(W)=0$, so gradings for W,M can both be calculated with respect to the trivialization of the canonical bundle of $\overline{M}$ (2.6). Now apply 1). $\square$

11.4. Obstructions to exact $ST^*L\hookrightarrow T^*N$, simply connected case.

Theorem 56. *Let L,N be closed simply connected n-manifolds, $n\geq 4$. For any exact contact hypersurface $ST^*L\hookrightarrow T^*N$, the following hold*

1. *$H^2(N)\to H^2(L)$ is injective;*
2. *$\pi_2(L)\to\pi_2(N)$ has finite cokernel;*
3. *if $H^2(N)\neq 0$ then $H_*(L)\cong H_*(W)$.*

Remark 57. *The maps in* (1), (2), (3) *are defined by the composites*

$\pi_2(L)\cong\pi_2(ST^*L)\to\pi_2(W)\to\pi_2(T^*N)\cong\pi_2(N)$,
$H^2(N)\cong H^2(T^*N)\to H^2(W)\to H^2(ST^*L)\cong H^2(L)$.

Using dim$(L)\geq 4$, we identify $\pi_1(L),\pi_2(L),H_1(L),H^2(L)$ with the analogous groups for $\partial W=ST^*L$ via the long exact sequence in homotopy for $S^{n-1}\to ST^*L\to L$ and the Gysin sequence $H^*(ST^*L)\to H^{*-(n-1)}(L)\to H^{*+1}(L)\to H^{*+1}(ST^*L)$.

Proof. Suppose by contradiction that $0\neq\eta\in H^2(N)$ vanishes in $H^2(L)\cong H^2(\partial W)$. By Corollary 70, $SH^*(T^*N)_\eta=0$. By Theorem 35, $SH^*(W)_{\eta|W}=0$. By Theorem 41, $SH^*_+(W)_{\eta|W}\cong H^{*+1}(W)\otimes\Lambda$ has finite rank. By Corollary 55, $SH^*_+(W)_{\eta|W}=BSH^*(ST^*L)_{\eta|\partial W}$. Using $\eta|_{\partial W}=0\in H^2(\partial W)$ and Corollary 42,

$$BSH^*(ST^*L)_{\eta|\partial W}\cong BSH^*(ST^*L)\otimes\Lambda\cong H_{n-*}(\mathcal{L}L,L)\otimes\Lambda.$$

This has infinite rank as $H_*(\mathcal{L}L)$ does [35], using L is closed with $\pi_1(L)=1$. So $SH^*_+(W)_{\eta|W}$ has both finite and infinite rank. Contradiction, proving (1).

The above shows that if $\tau\eta\neq 0$ in $H^1(\mathcal{L}N)$, then $\tau\eta|_{\partial W}\neq 0$ in $H^1(\mathcal{L}\partial W)\cong H^1(\mathcal{L}L)$ (using $\mathcal{L}N=\mathcal{L}_0N$ as $\pi_1(N)=1$). For $\pi_1(N)=1$, $\tau:H^2(N)\to H^1(\mathcal{L}N)$ is an isomorphism and $H^1(\mathcal{L}N)\cong\mathrm{Hom}(H_1(\mathcal{L}N),\mathbb{Z})\cong\mathrm{Hom}(\pi_2(N),\mathbb{Z})$. Thus

$$\mathrm{Hom}(\pi_2(N),\mathbb{Z})\to\mathrm{Hom}(\pi_2(L),\mathbb{Z}),\quad \tau\eta\mapsto\tau\eta|_{\partial W}$$

must be injective. Dualizing proves (2).

If $H^2(N) \neq 0$ then pick an $\eta \neq 0 \in H^2(N)$. As before, $SH^*(T^*N)_\eta = 0$ so $SH^*(W)_{\eta|W} = 0$. By (1) it restricts to $\eta|_{\partial W} \neq 0 \in H^2(ST^*L) \cong H^2(T^*L)$. By Corollary 70, $SH^*(T^*L)_{\eta|\partial W} = 0$. Using Theorem 41 and Corollary 55,

$$H^{*+1}(W) \otimes \Lambda \cong SH^*_+(W)_{\eta|W} \cong SH^*_+(DT^*L)_{\eta|\partial W} \cong H^{*+1}(L) \otimes \Lambda.$$

Passing to homology by universal coefficients, proves (3) (by 15.17 the claim holds for integral coefficients by defining Λ over $\mathbb{Z}$ instead of $\mathbb{K}$). $\square$

Conjecture. *For simply connected L, N, exact contact $ST^*L \subset T^*N$ always arise as the boundary of a Weinstein neighbourhood of an exact Lagrangian $L \hookrightarrow T^*N$.*

11.5. **Obstructions to exact $ST^*L \hookrightarrow T^*N$, general case.** The key to Theorem 56 was that $H_*(\mathcal{L}L)$ had infinite rank. This holds for closed L with $\pi_1(L) = 1$, since Sullivan [35] showed $H_*(\mathcal{L}L; \mathbb{Q})$ is infinite dimensional via rational homotopy theory. Sullivan's proof also works for nilpotent spaces. We do not know when the result holds for finite $\pi_1(L)$ (L need not be nilpotent: $\mathbb{R}P^2$ has $\pi_1 = \mathbb{Z}/2\mathbb{Z}$ but does not act nilpotently on $\pi_2 = \mathbb{Z}$). When $\pi_1(L)$ has infinitely many conjugacy classes (for example infinite abelian $\pi_1(L)$) the result holds because the connected components of $\mathcal{L}L$ are indexed by these conjugacy classes, so $\operatorname{rank} H_0(\mathcal{L}L) = \infty$.

Write $\pi : \widetilde{N} \to N$ for the universal cover, inducing $\pi^* : H^2(N) \to H^2(\widetilde{N})$.

Theorem 58. *Let L, N be closed n-manifolds, $n \geq 4$, with N of finite type (13.6). Let $ST^*L \hookrightarrow T^*N$ be an exact contact hypersurface. If $\operatorname{rank} H_*(\mathcal{L}_0 L) = \infty$ then:*

1. *$\pi_2(L) \to \pi_2(N)$ has finite cokernel.*
2. *if N is simply connected, $H^2(N) \to H^2(L)$ is injective;*
3. *$H^2(\widetilde{N}) \to H^2(\widetilde{L})$ and $H^2(N) \setminus \ker \pi^* \to H^2(L)$ are injective;*
4. *if L has finite type and $\operatorname{rank}(\pi_2 N) \neq 0$, then $H_*(L) \cong H_*(W)$.*

Now assume only that the full loop space $H_*(\mathcal{L}L)$ has infinite rank. Then (1), (2), (3) hold after replacing $\pi_2(L)$, $H^2(\widetilde{L})$ by $\pi_2(W)$, $H^2(\widetilde{W})$ respectively. Also if any one of (1) – (4) failed, then Corollary 60 (1) – (4) all hold and $\ker \pi^* = H^2(N)$ (so the image of the Hurewicz homomorphism $\pi_2(M) \to H_2(M)$ is torsion).

Remark 59. *The distinction (2), (3) is because we need $\tau\eta \neq 0 \in H^1(\mathcal{L}_0 N)$ in Corollary 70: $\tau : H^2(N) \to H^1(\mathcal{L}_0 N)$ is an isomorphism if $\pi_1(N) = 1$, but otherwise has kernel $\ker \pi^*$. We identify $H^2(\widetilde{N}) \cong \mathrm{Hom}(\pi_2(N), \mathbb{Z})$ with $\mathrm{im}\,(\tau) \subset H^1(\mathcal{L}_0 N)$ (6.6). The 2^{nd} part of (3) and (2) can be proved as in Theorem 56 using $\operatorname{rank} H_*(\mathcal{L}L) = \infty$. Note: (1) implies (2), (3). The 1^{st} part of (3) is dual to (1), the 2^{nd} part follows by*

$$\begin{tikzcd}
H^2(N) \arrow[r, "\tau"] \arrow[d] & \mathrm{Hom}(\pi_2(N), \mathbb{Z}) \arrow[d] \arrow[r, phantom, "\subset"] & H^1(\mathcal{L}_0 N) \\
H^2(L) \arrow[r, "\tau"] & \mathrm{Hom}(\pi_2(L), \mathbb{Z}) \arrow[r, phantom, "\subset"] & H^1(\mathcal{L}_0 L)
\end{tikzcd}$$

Proof. To prove (1) we twist by 2-forms $\widetilde{\eta} \neq 0 \in H^2(\widetilde{N})$ on the universal cover of N as explained in 6.6. By transgression, these give rise to 1-forms on the space $\mathcal{L}_0 N$ of contractible loops in N. By Theorem 20 there is a TQFT structure on the subgroup $SH^*_0(M)_{\widetilde{\eta}}$ of $SH^*(M)_{\widetilde{\eta}}$ generated by the contractible orbits.

Pass to universal covers and use transgressions (6.6) to identify the two rows:

$$\begin{tikzcd}
H^2(\widetilde{N}) \arrow[r] \arrow[d, equal] & H^2(\widetilde{W}) \arrow[r] \arrow[d, equal] & H^2(\widetilde{L}) \arrow[d, equal] \\
\mathrm{Hom}(\pi_2 N, \mathbb{Z}) \arrow[r] & \mathrm{Hom}(\pi_2 W, \mathbb{Z}) \arrow[r] & \mathrm{Hom}(\pi_2 L, \mathbb{Z})
\end{tikzcd}$$

Case 1: $\operatorname{rank} H_*(\mathcal{L}_0 L) = \infty$. Suppose by contradiction that $\widetilde{\eta}$ vanishes under this composition: $\widetilde{\eta}|_{\widetilde{L}} = 0 \in H^2(\widetilde{L})$. By [27], $SH^*_0(T^*N)_{\widetilde{\eta}} \cong H_{n-*}(\mathcal{L}_0 N)_{\widetilde{\eta}} = 0$ since N has finite type. As before, $SH^*_0(W)_{\widetilde{\eta}} = 0$, so $SH^*_{+,0}(W)_{\widetilde{\eta}} = H^{*+1}(W) \otimes \Lambda$ has finite rank. But, by independence of the filling, this group is isomorphic to

$$SH^*_{+,0}(DT^*L)_{\widetilde{\eta}|\widetilde{L}} = SH^*_{+,0}(DT^*L)_0 \cong H_{n-*}(\mathcal{L}_0 L, L) \otimes \Lambda,$$

which has infinite rank, contradiction. So the composition is injective, so dualizing gives (1). By Remark 59 it remains to prove (4). If $\operatorname{rank} \pi_2(N) \neq 0$, pick $\tau\widetilde{\eta} \neq 0$. By (2), also $\tau\eta|_{\widetilde{L}} \neq 0$, so since L has finite type, $SH^*_0(T^*L)_{\widetilde{\eta}|\widetilde{L}} = 0$. Thus

$$H^{*+1}(W) \otimes \Lambda \cong SH^*_{+,0}(W)_{\widetilde{\eta}} \cong SH^*_{+,0}(DT^*L)_{\widetilde{\eta}} \cong H^{*+1}(L) \otimes \Lambda.$$

Passing to homology by universal coefficients and using 15.17, proves (4).

Case 2: $\operatorname{rank} H_*(\mathcal{L}L) = \infty$. Assume by contradiction that $\widetilde{\eta}|_{\widetilde{W}} = 0 \in H^2(\widetilde{W})$. As before, $SH^*_0(W)_{\widetilde{\eta}} = 0$. Since $\widetilde{\eta}|_{\widetilde{W}} = 0$, this implies $SH^*_0(W) = 0$. By Theorem 35, $SH^*(W) = 0$ since the unit lies in $SH^*_0(W) = 0$. By Theorem 41, $SH^*_+(W) \cong H^{*+1}(W)$ has finite rank. But by Corollary 55 and Corollary 42,

$$SH^*_+(W) \cong BSH^*(ST^*L) \cong H_{n-*}(\mathcal{L}L, L),$$

which has infinite rank since $H_*(\mathcal{L}L)$ does. Contradiction. So $H^2(\widetilde{N}) \to H^2(\widetilde{W})$ is injective and dually $\pi_2(W) \to \pi_2(N)$ has finite cokernel. $\square$

11.6. **Pathological** L**.** Write $[S^1, L]$ for the set of free homotopy classes of maps $S^1 \to L$, equivalently: it is the set of conjugacy classes of $\pi_1(L)$.

Call L *pathological* if $[S^1, L] \neq 1$ is a finite set and $H_*(\mathcal{L}L)$ has finite rank. This is the only case when Theorem 58 may not apply.

Corollary 60. *Let L, N be closed n-manifolds, $n \geq 4$, with N of finite type and L pathological. Let $ST^*L \hookrightarrow T^*N$ be an exact contact embedding. Then either the results of Theorem 58 all hold, or else the following must hold:*

1. *$H_1(L) \to H_1(W)$ and $[S^1, L] \to [S^1, W]$ both vanish;*
2. *rank $H^{n+1}(W) = \#[S^1, L] - 1 \geq 1$;*
3. *if L is of finite type then $H^2(\widetilde{N}) \to H^2(\widetilde{L})$, $H^2(N) \backslash \ker \pi^* \to H^2(L)$ vanish;*
4. *if L is of finite type then the image of $\pi_2(L) \to \pi_2(N)$ is torsion.*

In particular, if W is Stein then the results of Theorem 58 hold.

Proof. For the last claim: $H^{n+1}(W) \neq 0$ by (2), but $H^{n+1}(W) = 0$ for Stein W^{2n}.

The connected components of $\mathcal{L}L$ are indexed by $[S^1, L]$. Filter $SH^*(DT^*L)$ by $[S^1, L]$ as in 15.13, and consider $i : [S^1, L] \cong [S^1, ST^*L] \to [S^1, W]$.

Suppose Theorem 58 fails. By Remark 59 some $\widetilde{\eta} \neq 0 \in H^2(\widetilde{N})$ has $\widetilde{\eta}|_{\widetilde{L}} = 0 \in H^2(\widetilde{L})$. By the proof of Theorem 58 $SH^*(W) = 0$ and, using Theorem 41,

$$H_0(\mathcal{L}L, L) \cong SH^n_+(DT^*L) \cong SH^n_+(W) \cong \bigoplus_{w \in [S^1, W]} SH^n_{+,w}(W) \cong H^{n+1}(W),$$

$$H_0(\mathcal{L}L, L) \cong \bigoplus_{0 \neq c \in [S^1, L]} H_0(\mathcal{L}_c L) \cong \bigoplus_{0 \neq c \in [S^1, L]} SH^n_{+,c}(DT^*L) \hookrightarrow \bigoplus_{0 \neq c \in [S^1, L]} SH^n_{+,i(c)}(W).$$

Now $H_0(\mathcal{L}_c L) = \mathbb{Z}$, so the two lines imply rank $H^{n+1}(W) = \#[S^1, L] - 1$. Also, $SH^n_{+,w}(W) \cong SH^n_w(W) = 0$ for $0 \neq w \in [S^1, W]$. So, since the second line respects summands, i must vanish (and similarly if we filter by H_1, 15.13), proving (1),(2).

Suppose $\widetilde{\eta}\in H^2(\widetilde{N})$ has $\widetilde{\eta}|_{\widetilde{L}} \neq 0 \in H^2(\widetilde{L})$. Then $SH_0^*(T^*N)_{\widetilde{\eta}}=0$, $SH_0^*(W)_{\widetilde{\eta}}=0$,

$$H^{*+1}(W) \otimes \Lambda \cong SH^*_{+,0}(W)_{\widetilde{\eta}} \cong SH^*_{+,0}(DT^*L)_{\widetilde{\eta}} \cong H_{n-*}(\mathcal{L}_0 L, L)_{\widetilde{\eta}}.$$

But for finite type L, $H_*(\mathcal{L}_0 L)_{\widetilde{\eta}} = 0$ since $\widetilde{\eta} \neq 0 \in H^2(\widetilde{L})$. Thus, $H_{n-*}(\mathcal{L}_0 L, L)_{\widetilde{\eta}} \cong H_{n-*-1}(L)$ by Corollary 42. So $H^{n+1}(W) \cong H_{-1}(L) = 0$ contradicts (2). Thus $H^2(\widetilde{N}) \to H^2(\widetilde{L})$ vanishes, proving the 1st part of (3). Taking transgressions and dualizing implies (4), which implies the 2nd part of (3) (Remark 59). □

11.7. Obstructions to exact contact embeddings.

Theorem 61. *Let M^{2n} be a Liouville domain with $c_1(M) = 0$, $n \geq 4$. Let L^n be a non-pathological closed manifold admitting an exact contact $i : ST^*L \hookrightarrow \overline{M}$. Then*

1. *If $SH^*(M)_\eta = 0$ then $i^*\eta \neq 0 \in H^2(L)$ and $\tau(i^*\eta) \neq 0 \in H^1(\mathcal{L}_0 L)$;*
2. *If $SH^*(M)_{\widetilde{\eta}} = 0$ then $i^*\widetilde{\eta} \neq 0 \in H^2(\widetilde{L})$;*
3. *If L is of finite type and $SH^*(M)_{\widetilde{\eta}} = 0$, then $H_*(L) \cong H_*(W)$.*

For pathological L: if (1) or (2) fails, then Corollary 60(1)(2) hold, W is not Stein. For finite type pathological L: $SH^*(M)_\eta=0$, $SH^*(M)_{\widetilde{\eta}}=0$ imply $\tau(i^*\eta)=0$, $i^*\widetilde{\eta}=0$.

Corollary 62. *Under the assumptions of 61, if $SH^*(M)=0$ or $SH^*(W)=0$, then:*

- *L must be pathological, W is not Stein;*
- *$H_1(L) \to H_1(W)$ and $[S^1, L] \to [S^1, W]$ both vanish;*
- *rank $H^{n+1}(W) = \#[S^1, L] - 1 \geq 1$.*

Example. Corollary 62 applies to subcritical Stein manifolds M with $c_1(M) = 0$, since $SH^*(M) = 0$ by Cieliebak [10]. This mildly strengthens [12, Cor. 1.18]

12. DISPLACEABILITY OF ∂M AND RABINOWITZ FLOER THEORY

12.1. Rabinowitz Floer cohomology.

The Lagrange multiplier analogue of SH^* is *Rabinowitz Floer cohomology*, due to Cieliebak-Frauenfelder [11]. It involves changing the action A_H by rescaling H by a *Lagrange multiplier* variable $\lambda \in \mathbb{R}$:

$$L_H(x, \lambda) = -\int x^*\theta + \lambda \int_0^1 H(x(t))\, dt.$$

This tweak has a considerable effect: it forces *all* critical points of the new action functional L_H to lie *in the boundary* ∂M. Choosing $X_H = \mathcal{R}$ on ∂M, the critical points are the solutions of $\dot{x} = \lambda \mathcal{R}(x)$ and they have action $L_H = -\lambda$. So, after reparametrizing $y(t) = x(t/|\lambda|)$, these critical points are precisely: the Reeb orbits of period $\lambda > 0$, the constant loops in ∂M ($\lambda = 0$), and the negative Reeb orbits of period $-\lambda > 0$. The chain complex generated by the critical points of L_H has a differential defined by counting negative gradient flowlines of L_H. The resulting cohomology is called *Rabinowitz Floer cohomology* $RFH^*(M)$ (see [11]).

Lemma 63 (Cieliebak-Frauenfelder-Oancea, [12])**.** *There is a long exact sequence*

$$\cdots \to SH_{2n-*}(M) \to SH^*(M) \to RFH^*(M) \to SH_{2n-*-1}(M) \to \cdots$$

where the map $SH_{2n-*}(M) \to SH^*(M)$ is the composite

$$SH_{2n-*}(M) \xrightarrow{c_*} H_{2n-*}(M) \xrightarrow{\text{Poincaré}} H^*(M, \partial M) \xrightarrow{\text{inclusion}^*} H^*(M) \xrightarrow{c^*} SH^*(M).$$

Remark 64. *The Lemma is written in our conventions. Comparison: conventions in [12] dictate $SH_*(DT^*N) \cong H_*(\mathcal{L}N)$ instead of $SH^{n-*}(DT^*N) \cong H_*(\mathcal{L}N)$.*

12.2. Vanishing theorem.

Theorem 65. *$RFH^*(M) = 0$ iff $SH^*(M) = 0$ iff $SH_*(M) = 0$.*
If one among SH_0, SH^0, RFH^0 is 0 (even just on the component $\mathcal{L}_0\overline{M}$ of contractible loops, 15.13), then all SH_*, SH^*, RFH^* are 0. Here we used $\mathbb{Z}$-gradings if $c_1 = 0$ (or $c_1(\pi_2 M) = 0$ if we restrict to $\mathcal{L}_0\overline{M}$) otherwise use $\mathbb{Z}/2\mathbb{Z}$-gradings.

Proof. By Theorem 38, $SH^*(M) = 0$ iff $SH_*(M) = 0$ iff one of them vanishes in degree 0 (where unit/counit lie). By Lemma 63, if $SH^*(M) = SH_*(M) = 0$ then $RFH^*(M) = 0$. Conversely, suppose $RFH^0(M) = 0$ (even restricted to contractible loops, since the long exact sequence of Lemma 63 respects filtrations 15.13). By Theorems 35, 38 it suffices to show that the unit $e \in SH^0(M)$ vanishes.

Case 1. Suppose we have a $\mathbb{Z}$-grading. Lemma 63 yields the exact sequence $SH_{2n}(M) \to SH^0(M) \to RFH^0(M)$, whose first map is the composite

$$SH_{2n}(M) \to H_{2n}(M) \to H^0(M, \partial M) \to H^0(M) \to SH^0(M).$$

Now M is a $2n$-manifold with non-empty boundary so $H_{2n}(M) = 0$. Thus the above composite is zero, so in the above exact sequence, $SH^0(M) \hookrightarrow RFH^0(M)$ is injective. Thus $RFH^0(M) = 0$ implies $SH^0(M) = 0$ so $e = 0$.

Case 2. Suppose we have a $\mathbb{Z}/2\mathbb{Z}$-grading. By Lemma 63, e is in the image of $SH_{\text{even}}(M) \to SH^{\text{even}}(M)$ since it maps to zero via $SH^{\text{even}}(M) \to RFH^{\text{even}}(M) = 0$. The $H^0(M)$ summands of the image of $SH_{\text{even}}(M) \to H_{\text{even}}(M) \to H^{\text{even}}(M, \partial M) \to H^{\text{even}}(M)$ must vanish since they factor through $H_{2n}(M) = 0$. So since e is in the image of $SH_{\text{even}}(M) \to SH^{\text{even}}(M)$, $e = c^*(z)$ for some $z \in H^*(M) \setminus H^0(M)$. But c^* is a TQFT map and $z^n = 0$ for degree reasons, so $0 = c^*(z^n) = c^*(z)^n = e^n = e$. □

12.3. Displaceability of ∂M implies vanishing of $SH^*(M)$.

An exact symplectic manifold $(X^{2n}, d\theta)$ is *convex* if it is connected, without boundary, admitting an exhaustion $X = \cup_k X_k$ by compacts $X_k \subset X_{k+1}$, such that the Liouville vector field Z defined by $i_Z d\theta = \theta$ points strictly outwards along ∂X_k, Z is complete (its flow is defined for all time) and $Z \neq 0$ outside of some X_k. **Example.** $X = \overline{M}$.

Theorem 66. *If M admits an exact embedding into an exact convex symplectic manifold X, such that ∂M is displaceable by a compactly supported Hamiltonian flow in X, then: $SH^*(M) = 0$, there are no closed exact Lagrangians in M, and the Arnol'd chord conjecture holds for any Legendrian arising as in 3.1.*

Proof. If $\partial M \subset X$ is displaceable then $RFH^*(\partial M \subset X) = 0$ by Cieliebak-Frauenfelder [11]. But RFH^* only depends on the filling M of ∂M, so $RFH^*(M) = 0$. By Theorem 65, $SH^*(M) = 0$. If there were a closed exact $L \subset M$ then for $W \cong DT^*L$ as in Example 28 the restriction $0 = SH^*(M) \to SH^*(W) \cong H_{n-*}(\mathcal{L}L)$ implies $H_*(\mathcal{L}L) = 0$ by Theorem 35, absurd. For the last claim use Theorems 40, 44. □

Example. If $ST^*L \hookrightarrow X$ is exact contact such that ST^*L is displaceable, $c_1(X) = 0$, $n \geq 4$, then $SH^*(W) = 0$ so Corollary 62 holds. So for non-pathological L, ST^*L is never displaceable (mildly strengthens Thm 1.17 of [12]).

13. String topology

For closed n-manifolds N, Viterbo [36] proved $SH^*(T^*N) \cong H_{n-*}(\mathcal{L}N)$, and there are now other approaches by Abbondandolo-Schwarz [1] and Salamon-Weber [30]. We use approach [1] since we used it in [27] to prove $SH^*(T^*N)_\eta \cong H_{n-*}(\mathcal{L}N)_\eta$. We claim that these isomorphisms respect the TQFT. We write $H_*(\mathcal{L}N)_\eta$ to keep track of twistings by $\tau\eta$ when present. We tacitly assume that N^n is orientable, although the non-orientable case also holds if $\mathrm{char}(\mathbb{K}) = 2$.

13.1. **The Abbondandolo-Schwarz construction.** We first identify $H^*(\mathcal{L}N)_\eta$ with the Morse cohomology $MH^*(\mathcal{L}N)_\eta$ of the Lagrangian action functional

$$\mathcal{E}(\gamma) = \textstyle\int_0^1 L(\gamma, \dot\gamma)\, dt,$$

for a suitable function $L : TN \to \mathbb{R}$ which is quadratic in the fibres, for example the Lagrangians $L(q, v) = \frac{1}{2}|v|^2 - U(q)$ from classical mechanics.

We replace $\mathcal{L}N = C^\infty(S^1, N)$ with the homotopy equivalent space $W^{1,2}(S^1, N)$, and one proves that Morse homology for $\mathcal{E}$ on this space is well-defined for most L.

The choice of L determines the choice of a quadratic Hamiltonian on T^*N,

$$H(q, p) = \max_{v\in T_qN} (p \cdot v - L(q, v)).$$

The convexity of L implies that the maximum is achieved at precisely one point $p = d_vL$. This defines a fiber-preserving diffeomorphism, the Legendre transform

$$\mathfrak{L} : TN \to T^*N, (q, v) \mapsto (q, d_vL(q, v)).$$

This relates the dynamics of H on T^*N with that of $\mathcal{E}$ on $\mathcal{L}N$: $\gamma \in \mathrm{Crit}(\mathcal{E})$ is a critical point of $\mathcal{E}$ in $\mathcal{L}N$ if and only if $x = \mathfrak{L}(\gamma, \dot\gamma) \in \mathrm{Crit}(A_H)$ is a 1-periodic orbit of X_H in T^*N. One can even relate the indices and the action values of $\mathcal{E}$ and A_H.

The isomorphism $H_*(\mathcal{L}N)_\eta \to SH^{n-*}(T^*N)_\eta$ is induced by a chain map

$$\varphi : MC_*(\mathcal{E})_\eta \to SC^{n-*}(H)_\eta, \quad \varphi(\gamma) = \sum_{v\#u\in\mathcal{M}_0^+(\gamma,x)} \epsilon_{v\#u}\, t^{-\tau\eta[v]+\tau\eta[\pi u]}\, x$$

where $\epsilon_{v\#u} \in \{\pm1\}$ are orientation signs and where $\mathcal{M}^+(\gamma, x)$ consists of pairs $v\#u$, with $v : (-\infty, 0] \to \mathcal{L}N$ a $-\nabla\mathcal{E}$ flowline converging to $\gamma \in \mathrm{Crit}(\mathcal{E})$, and $u : (-\infty, 0]\times S^1 \to T^*N$ solves Floer's equation and converges to $x \in \mathrm{Crit}(A_H)$, with Lagrangian boundary condition $\pi\circ u(0, t) = v(0, t)$ ($\pi : T^*N \to N$ is the projection).

As usual, one makes a C^2-small time-dependent perturbation of L (hence H), but we keep the notation simple. Appendix 17 explains why we can use a quadratic Hamiltonian instead of taking a direct limit over linear Hamiltonians.

13.2. **TQFT structure on $H_*(\mathcal{L}N)$.** Given a graph S' as in 15.20, use the Morse function $\mathcal{E}_i(\gamma) = \int_0^1 L_i(\gamma, \dot\gamma)\, dt$ for edge e_i for a generic $L = L_i$ as above. This yields

$$\psi_{S'} : MH_*(\mathcal{L}N)_\eta^{\otimes p} \to MH_*(\mathcal{L}N)_\eta^{\otimes q}.$$

For example $\psi_{Q'} : MH_*(\mathcal{L}N)_\eta^{\otimes 2} \to MH_*(\mathcal{L}N)_\eta$ is a product (Figure 12). Identifying $MH_*(\mathcal{L}N) \cong H_*(\mathcal{L}N)$, these Morse operations define a TQFT on $H_*(\mathcal{L}N)$. A string topology focused description of these can be found in Cohen-Schwarz [13].

13.3. **The Chas-Sullivan loop product.** Let $ev : \mathcal{L}N \to N$ be evaluation at 0. Let $\sigma : \Delta^a \to \mathcal{L}N$ and $\tau : \Delta^b \to \mathcal{L}N$ be two singular chains, think of them as a, b-dimensional families of loops. When two loops from those two families happen to have the same base-point, we can form a "figure 8" loop. Let $E = \{(s, t) \in \Delta^a\times\Delta^b : ev(\sigma(s)) = ev(\tau(t))\}$ parametrize those match-ups and let $E \to \mathcal{L}N$ output those figure-8 loops. Now $E = (ev \times ev)^{-1}(\text{diagonal})$ via $ev \times ev : \Delta^a \times \Delta^b \to N \times N$. So E is a manifold when the two families of basepoints, $ev(\sigma)$ and $ev(\tau)$, intersect transversely in N, and $\dim(E) = a + b - n$. Thus $E \to \mathcal{L}N$ defines a chain in $H_{a+b-n}(\mathcal{L}N)$: the *loop product* $\sigma\cdot\tau$ (see [2] for details).

Lemma 67 (Abbondandolo-Schwarz [2])**.** *The isomorphism* $\varphi : MH_*(\mathcal{L}N) \to SH^{n-*}(T^*N)$ *preserves the product structure, so* $\varphi \circ \psi_{Q'} = \psi_P \circ \varphi^{\otimes 2}$*. The product* $\psi_{Q'}$ *on* $MH_*(\mathcal{L}N)$ *corresponds to the* Chas-Sullivan loop product [9] *on* $H_*(\mathcal{L}N)$*.*



FIGURE 3. Dark circles are critical points of $\mathcal{E}$, dark rectangles are Hamiltonian orbits. POP are pairs of pants, Floer are Floer trajectories. On the left the configurations counted by φ, $\psi_{Q'}$, ψ_P.

Sketch Proof. Observe Figure 3. Consider the 1-dimensional family of solutions corresponding to the configurations in the middle column (ℓ is a free parameter). On the right are the limiting configurations as $\ell \to 0$ or ∞. The middle two pictures in the last column are the same. So the maps counting the top and bottom configurations are chain homotopic. These are respectively $\varphi \circ \psi_{Q'}$ and $\psi_P \circ \varphi^{\otimes 2}$. □

13.4. φ **is a TQFT isomorphism.** To S we associated a graph S' (15.19). Let $\overline{S'}$ be the graph obtained from S' by switching the orientations of the edges (p, q get interchanged). **Example.** for P, Q of 15.18 we get the P', Q' of 15.19 and $\overline{P'} = Q'$.

Theorem 68. *The isomorphism* $\varphi : H_*(\mathcal{L}N)_\eta \cong MH_*(\mathcal{L}N)_\eta \to SH^{n-*}(T^*N)_\eta$ *preserves the (possibly twisted) TQFT structures:* $\varphi^{\otimes p} \circ \psi_{\overline{S'}} = \psi_S \circ \varphi^{\otimes q}$*.*
In particular, $\varphi(c_*[N]) = e$ *where* $c : N \to \mathcal{L}N$ *is inclusion of constants.*

Proof. Recall Z, C, P, Q from Corollary 99. For $S = Z$, ψ_Z and $\psi_{\overline{Z'}}$ are the identity, so there is nothing to prove. In fact, the proof for cylinders at the chain level is the same as the proof that φ is a chain map respecting action filtrations.

For $S = C$, one can argue indirectly that Floer solutions on C are finite energy cylinders by Lemma 96, which reduces us to the case of cylinders. More directly, by the compatibility of the unit with direct limits (Theorem 98) and using Appendix 17, or alternatively arguing by action restriction, we can assume the Hamiltonian is H^δ (4). We can assume H^δ has just one minimum x_0 on $N \subset T^*N$, which is the action minimizer. Thus the only isolated solution $u : S \to \overline{M}$ is the constant at x_0. So in the notation of 13.1, $\mathcal{M}^+(x_0, y) = \{v\#u\}$ where $u \equiv v \equiv x_0$ are constant, $x_0 = \min H = \max L|_{c(N)}$. Thus, $\varphi^{-1}(e)$ is the maximum of L on N, which in the ordinary cohomology $H^*(N)$ is the unit 1. Dually, this is the fundamental class $[N] \in H_n(N)$, so in $H_n(\mathcal{L}N)$ it corresponds to the image $c_*[N]$.

For $S = P$, the relation $\varphi \circ \psi_{Q'} = \psi_P \circ \varphi^{\otimes 2}$ is Lemma 67. The techniques of [2] can also be applied to the pair of pants $S = Q$, so $\varphi^{\otimes 2} \circ \psi_{P'} = \psi_Q \circ \varphi$.

For general S, decompose S into copies of C, Z, P, Q as in Corollary 99. Thus ψ_S is a composition of operations $\psi_C, \psi_Z, \psi_P, \psi_Q$ and $\psi_{\overline{S'}}$ is the corresponding composition of operations obtained by replacing C, Z, P, Q by $\overline{C'}, \overline{Z'}, \overline{P'}, \overline{Q'}$. Thus the general relation follows since it holds C, Z, P, Q.

For the twisted case, φ was constructed in [27]. For the argument above to hold in the twisted case, it suffices that the weights are locally constant on the TQFT moduli spaces and the $\mathcal{M}_0^+(\gamma, x)$ moduli spaces. The former is done in Theorem 15 and the latter is done in [27]. $\square$

Corollary 69. *1) Via φ, the map $c^* : H^*(T^*N) \otimes \Lambda \to SH^*(T^*N)_\eta$ becomes the map $c_* : H_*(N) \otimes \Lambda \to H_*(\mathcal{L}N)_\eta$, induced by the inclusion of constants $c : N \to \mathcal{L}N$. This c_* is a TQFT map, using the intersection product on $H_*(N)$ with unit $[N]$, and using the twisted loop product on $H_*(\mathcal{L}N)_\eta$ with unit $c_*[N]$.*

2) For a closed exact Lagrangian submanifold $L \subset T^*N$, Theorem 32 applied Example 28 yields the commutative TQFT diagram of transfer maps

$$\begin{tikzcd}
H_*(\mathcal{L}L)_{\eta|L} & H_*(\mathcal{L}N)_\eta \arrow[l] \\
H_*(L) \otimes \Lambda \arrow[u, "c_*"] & H_*(N) \otimes \Lambda \arrow[l] \arrow[u, "c_*"]
\end{tikzcd}$$

Remark. The untwisted diagram (without TQFT) is due to Viterbo [37].

Proof. Claim 1) identifies $H^*(T^*N) \cong H^*(N)$ and applies Poincaré duality. The dual of $c^*(1) = e$ (Theorem 95) shows that $[N]$ and $c_*[N]$ are units. The *intersection product* on $H_*(N)$ is the dual of the cup product [8] (take the exterior product $H_a(N) \otimes H_b(N) \to H_{a+b}(N \times N)$ and push forward via the diagonal map $\Delta_! : H_{a+b}(N \times N) \to H_{a+b-n}(N)$, where $\Delta_!$ is Poincaré dual to the pullback Δ^*). $\square$

13.5. **The based loop space.** The wrapped analogue of $SH^*(T^*N) \cong H_{n-*}(\mathcal{L}N)$ is Abbondandolo-Schwarz's ring isomorphism [1] for a fibre $T^*_{q_0}N \subset T^*N$:

$$HW^*(T^*_{q_0}N) \cong H_{n-*}(\Omega N),$$

using the Pontrjagin product on $H_*(\Omega N)$ induced by concatenation of based loops. As in Theorem 68, this respects the TQFT and the twisted analogue holds.

13.6. **Vanishing of the Novikov homology of the free loop space.** A space N has *finite type* if $\pi_m(N)$ is finitely generated for each $m \geq 2$.

Examples. 1) simply connected closed manifolds, and
2) closed manifolds with trivial π_1 action on higher homotopy groups.

For closed N of finite type, the Novikov homology $H_*(\mathcal{L}_0N; \underline{\Lambda}_{\tau\eta}) = 0$ for $\tau\eta \neq 0 \in H^1(\mathcal{L}_0N)$ [27] ($\mathcal{L}_0N$ is the component of contractible loops). We extend it to $\mathcal{L}N$:

Corollary 70. *For closed manifolds N of finite type, and any $\tau\eta \neq 0 \in H^1(\mathcal{L}_0N)$,*

$$SH^*(T^*N)_\eta \cong H_{n-*}(\mathcal{L}N; \underline{\Lambda}_{\tau\eta}) = 0.$$

Proof. By Theorem 68, $SH^*(T^*N)_\eta \cong H_{n-*}(\mathcal{L}N)_\eta$ is a ring with unit $c_*[N] \in H_n(\mathcal{L}_0N)_\eta$. But $H_*(\mathcal{L}_0N)_\eta = 0$ by [27]. So the claim follows by Theorem 35. $\square$

Remark. $\tau\eta \neq 0 \in H^1(\mathcal{L}_0N)$ iff $\pi^*\eta \neq 0 \in H^2(\widetilde{N})$ for the universal cover $\pi : \widetilde{N} \to N$.

14. The c^* maps preserve the TQFT structure

14.1. **PSS map.** Theorem 14 is the analogue of the Piunikhin-Salamon-Schwarz [26] ring isomorphism $FH^*(M,H) \to QH^*(M)$ for (suitable) *closed* symplectic manifolds M. It turns the pair of pants product on Floer cohomology into the quantum cup product. We need to briefly survey their proof.



$\phi : CF^*(H) \to CM^*(f)$



$\psi : CM^*(f) \to CF^*(H)$

Pick a generic Morse function f and a Hamiltonian H on M. Pick a homotopy H_s interpolating 0 and H. The PSS map $\phi : CF^*(H) \to CM^*(f)$ on generators is $\phi(y) = Nx + \cdots$ where N is an oriented count of isolated *spiked discs* converging to x, y. Spiked discs are maps $u : \mathbb{C} \to M$ such that $u(e^{2\pi(s+it)})$ satisfies Floer's continuation equation for H_s, converging at $s = \infty$ to a 1-periodic orbit y of H and at $s = -\infty$ to a point $u(0)$ in the unstable manifold $W^u(x,f)$ of f. The "spike" is the $-\nabla f$ flow line connecting x to $u(0)$. An inverse $\psi : CM^*(f) \to CF^*(H)$ (up to chain homotopy) is defined by counting spiked discs flowing in the reverse direction: $u(e^{2\pi(-s+it)})$ is an isolated Floer continuation solution for H_{-s} converging at $s = -\infty$ to an orbit y of H and at $s = \infty$ to a point $u(0)$ in a stable manifold $W^s(x,f)$.



Figure 4. Counting the solutions defining $\phi \circ \psi$ and $\psi \circ \phi$.

To show that ϕ, ψ are inverses up to chain homotopy, consider $\phi \circ \psi$ and $\psi \circ \phi$. After gluing solutions, these count configurations on the left in Figure 4. Up to chain homotopy, the maps don't change if we homotope respectively H and ℓ to zero. So we count the configurations on the right. In the first case, for dimensional reasons the J-holomorphic sphere in the figure is constant. So we count isolated Morse continuation solutions for a constant homotopy f, which are constants (otherwise there is a 1-family of reparametrized solutions via $s \to s+\text{constant}$). So $\phi\circ\psi \simeq \text{id}$ are chain homotopic. In the second case, we used the gluing theorem for J-holomorphic curves to obtain a cylinder. Up to chain homotopy, we can homotope H_s to an s-independent H. So we count isolated Floer continuation solutions for a constant homotopy H, which again must be constants. So $\psi \circ \phi \simeq \text{id}$.

To prove that ϕ, ψ respect the ring structure observe Figure 5. We glue spiked discs to cap off, respectively, Floer and quantum-Morse product solutions (the quantum product replaces the vertex of the Morse-product graph P' of 15.20 with a J-holomorphic sphere). Up to chain homotopy, we can homotope respectively H and ℓ_1, ℓ_2, ℓ_3 to zero, reducing to the configurations on the right. In the second



FIGURE 5. Capping off Floer and Morse product solutions.

case, we used the gluing theorem for J-holomorphic curves to get a pair of pants Floer solution. So up to chain homotopy, we obtain the products on QH^* and FH^* respectively. A similar argument works for all (quantum) TQFT operations.

14.2. **Proof for Liouville domains.** For Liouville domains M the compactness of the moduli spaces fails, otherwise ϕ, ψ would always induce $SH^*(M) \cong H^*(M)$ which is false. Compactness fails for ϕ since it uses a non-monotone homotopy H_s. $\psi : CM^*(f) \to SC^*(H)$ is well-defined as we can pick H_{-s} monotone (Lemma 81).

Lemma 71. *$\psi : HM^*(f) \to SH^*(H^\delta)$ equals $c^* : H^*(M) \cong SH^*(H^\delta)$.*

Proof. H^δ is C^2-small Morse (section 4) so the Floer solutions counted by ψ are time-independent. Now ψ changes by a chain homotopy if we homotope f, H_s to H, H. So we end up counting isolated Morse continuation solutions for the constant homotopy H, which must be constant solutions. So $\psi \simeq c^*$ are chain homotopic. □

Goal: To prove Theorem 14 by the methods of 14.1 it remains to construct a well-defined map $\phi : SH^*(H^\delta) \to H^*(M)$ inverse to ψ. We will do this in 14.4.

14.3. **Floer trajectories converging to broken Morse trajectories.** We recall the arguments [31] to prove $FH^*(H) \cong H^*(M)$ for C^2-small H on (suitable) closed symplectic manifolds M. Let $S^1_T = \mathbb{R}/T\mathbb{Z}$. We will be considering T-periodic Floer theory, as opposed to the 1-periodic theory, and we will study the limit $T \to 0$.

Consider solutions $u : \mathbb{R} \times S^1_T \to \mathbb{R}^{2n}$ to Floer's local equation $F_T(u) = 0$, where $F_T : W^{1,2}(\mathbb{R} \times S^1_T; \mathbb{R}^{2n}) \to L^2(\mathbb{R} \times S^1_T; \mathbb{R}^{2n})$ is the local Floer operator defined by

$$F_T(u) = \partial_s u + J\partial_t u + (S + A)u,$$

where J is standard, $S = S(s,t)$ and $A = A(s,t)$ are continuous matrix valued functions on $\mathbb{R} \times S^1_T$ which are respectively symmetric and antisymmetric, and which converge to $S_\pm$ and $A_\pm = 0$ as $s \to \pm\infty$.

Claim 1. *The solutions u of $F_T(u) = 0$ are time-independent whenever $\|S(s)\|_{L^2(dt)} + \|A(s)\|_{L^2(dt)} \leq c < 1/T$ for some constant c.*

Sketch Proof. [31, Prop.4.2] The averaged solution $\overline{u}(s) = (\int_0^T u(s,t)\,dt)/T$ is a solution $F_0(\overline{u}) = 0$ of the local Morse operator

$$F_0(\overline{u}) = \partial_s \overline{u} + (S + A)\overline{u},$$

and then one checks that $u(s,t) \equiv \overline{u}(s)$ when S, A are small in the above sense. □

Assume in addition that $S_\pm$ are non-singular, which is the local equivalent of the non-degeneracy condition of orbits in Floer theory.

Claim 2. *F_T is onto iff F_0 is onto.*
Sketch proof. [31, Cor.4.3] This follows because they have the same Fredholm index, and by Claim 1 they have isomorphic kernels. □

The local result can now be applied globally. Pick an ω-compatible almost complex structure on the closed symplectic manifold (M,ω) such that $\omega(\cdot, J\cdot)$ is a Morse-Smale metric for a given Morse function $H : M \to \mathbb{R}$.

Claim 3. *For small $T > 0$, every finite energy Floer trajectory $u : \mathbb{R} \times S^1_T \to M$ is time-independent and regular (the linearization of the Floer operator at u is onto).*
Sketch proof. [31, Thm.7.3] When u is a time-independent Floer trajectory, i.e. a $-\nabla H$ flow, we reduce to the above local case by picking an orthonormal frame: F_0 is onto by the Morse-Smale condition, so F_T is onto, so u is regular.

To prove that u must be time-independent we need a Gromov compactness trick. By contradiction suppose there are finite energy Floer solutions $u_n : \mathbb{R} \times S^1_{T_n} \to M$ with periods $T_n \to 0$, which are not time-independent, connecting two given critical points x, y of H (we assume T_n is so small that there are no T_n-periodic Hamiltonian orbits, this uses the fact that H is Morse). The energy

$$E(u_n) = \tfrac{1}{T_n} \int_{-\infty}^{\infty} \int_0^{T_n} |\partial_s u_n|^2 \, dt \, ds = H(y) - H(x)$$

is bounded, therefore by Gromov compactness, a subsequence of the u_n must converge to some broken Morse trajectory $v_1 \# \cdots \# v_m$ for H, where m is bounded by the difference of the Morse indices of x, y. For simplicity, assume the u_n are isolated Floer solutions. Then $m = 1$, and $v = v_1$ is an isolated Morse trajectory, i.e. a time-independent isolated Floer trajectory.

We claim that for large n, $u_n(s + s_n', t) \equiv v(s)$ coincide for some $s_n' \in \mathbb{R}$. This would contradict the assumption that u_n is not time-independent. So suppose by contradiction that u_n and v never coincide in this way.

If there are integers C_n with $C_n T_n = T$, with T so small that F_T is onto at v by the first part of the proof, then $F_T(u_n) = 0$ and $u_n \to v$ contradicts the fact that v is an isolated solution of F_T. In general, pick integers C_n with $C_n T_n \to T$ and apply the same argument uniformly in an interval around T. □

Claim 4. *Given a Morse function $H : M \to \mathbb{R}$ and a small enough $T > 0$, all Floer trajectories for $T \cdot H$ are time-independent and regular.*
Proof. The T-periodic and 1-periodic Floer theories are isomorphic, by sending a T-periodic Floer trajectory $u(s,t)$ for H to the 1-periodic Floer trajectory $u(sT, tT)$ for $T \cdot H$. By Claim 3, for $T \ll 0$, the T-periodic Floer trajectories for H are time-independent and regular, so the same is true for the 1-periodic ones for $T \cdot H$. □

We can apply the same arguments to Floer continuation solutions $u : \mathbb{R} \times S^1_T \to M$ and Morse continuation solutions for homotopies $H_s : M \to \mathbb{R}$ connecting two Morse functions and picking J_s so that $g_s = \omega(\cdot, J_s \cdot)$ is Morse-Smale for H_s. Thus,

Claim 5. *For small $T > 0$, all Floer continuation solutions for $T \cdot H_s$ are time-independent and regular.*

14.4. **Non-monotone homotopies of small Hamiltonians.** We now achieve the "Goal" of 14.2. Let M be a Liouville domain. From now on $H = H^\delta$ (Sec. 4).

Pick the Morse function f and the homotopy H_s from 0 to H to be of the form $f = f(R)$, $H_s = h_s(R)$ on the collar, with $f'(R) > 0$ and $h'_s(R) > 0$ (unless $H_s \equiv 0$). This ensures that $-\nabla f$ and $-\nabla H_s$ are inward-pointing on the collar.

Definition 72. *Let $v : (-\infty, 0] \to \overline{M}$ and $u : \mathbb{C} \to \overline{M}$. Then $v\#u$ is a* ϕ-solution *from $x \in Crit(f)$ to $y(t) \equiv y \in Crit(H)$ if v is a $-\nabla f$ flow-line from x to $u(0)$, and $u(e^{2\pi(s+it)})$ is a Floer continuation solution for H_s converging to $y(t)$ at $s = \infty$. We write $\mathcal{M}^{\phi}(x, y)$ for the moduli space of ϕ-solutions. We call it a* T-periodic ϕ-solution *if we use $u(e^{2\pi(s+it)/T})$ and parametrize t by $\mathbb{R}/T\mathbb{Z}$ instead of $\mathbb{R}/\mathbb{Z}$.*

Lemma 73. *Fix a compact $M' \subset \overline{M}$ containing M. For small enough $T > 0$ all ϕ-solutions for $T \cdot H_s$ lying in M' are time-independent, regular and lie in M (provided the almost complex structures J,J_s induce Morse-Smale metrics for H,H_s.)*

Proof. Apply the Gromov compactness trick of 14.3: suppose by contradiction there are T_n-periodic ϕ-solutions $v_n\#u_n \in \mathcal{M}^{\phi}(x, y)$ lying in M', with $T_n \to 0$, such that u_n is not time-independent. The $v_n\#u_n$ have bounded energy

$$\begin{aligned} E(v_n) + E(u_n) &= (f(x) - f(u(0))) + (H(u(0)) - H(y)) \\ &\leq f(x) - H(y) + \max(f|_M) + \max(H|_M) \end{aligned}$$

because $u(0)$ must lie in M (since $-\nabla f$ is inward-pointing on the collar). So by Gromov compactness we can extract a subsequence converging to a broken Morse solution. Just like in Claim 3 of 14.3, we can then conclude that the u_n are time-independent for large n, which is a contradiction. Thus for small enough $T > 0$, all T-periodic ϕ-solutions lying in M' are time-independent and regular.

For time-independent ϕ-solutions, since the ends lie in M, the flow-line cannot exit M otherwise $-\nabla f$ or $-\nabla H_s$ would point outward at some point on the collar. The result follows by a rescaling trick like in Claim 4 of 14.3. □

Similarly define ψ-solutions, $\mathcal{M}^{\psi}(y, x)$ for f, H_{-s} (14.1), with H_{-s} monotone.

Lemma 74. *All ψ-solutions are contained in M. Moreover, for small $T > 0$, all ψ-solutions for $T \cdot H_{-s}$ are time-independent, regular, and contained in M.*

Definition 75. *Replace H_s by $T \cdot H_s$ so that the previous two lemmas hold. Define $\phi : SC^*(H) \to MC^*(f)$ and $\psi : MC^*(f) \to SC^*(H)$ by counting only the isolated ϕ, ψ-solutions* ***which lie in* M** *(ignore the ϕ-solutions outside M').*

Lemma 76. *ϕ and ψ are chain maps.*

Proof. Consider a 1-dimensional connected component $\mathcal{M} \subset \mathcal{M}_1^{\psi}(y, x)$. The boundary consists of broken ψ-solutions, where a Floer or a Morse trajectory breaks off at one of the two ends. The count of these broken solutions proves that $d_{\mathrm{Floer}} \circ \psi - \psi \circ d_{\mathrm{Morse}} = 0$. The same proof works for ϕ by Lemma 77, since we can ignore those $\mathcal{M} \subset \mathcal{M}_1^{\phi}(x, y)$ which contain a ϕ-solution which escapes M'. □

Lemma 77. *Let $\mathcal{M}$ be a connected component of $\mathcal{M}^{\phi}(x, y)$. Then $\mathcal{M}$ consists either entirely of solutions lying in M or entirely of solutions which escape M'.*

Proof. By Lemma 73, ϕ-solutions lying in M' must lie in M, so they cannot be C^0-close to a ϕ-solution which escapes M'. □

Theorem 78. *ψ, ϕ are inverse to each other up to chain homotopy. In particular, $\phi : SH^*(H^{\delta}) \to H^*(M)$ is an inverse for $\psi = c^* : H^*(M) \to SH^*(H^{\delta})$.*

Proof. Run the argument of 14.1, using that all J-holomorphic spheres are constant since the symplectic form is exact. The proof of Figure 4 involves the connected 1-dimensional components $\mathcal{M}$ of the moduli space $\mathcal{M}' = \bigcup_{c\in[0,1]} \mathcal{M}^{\phi\circ\psi}(x,x';f,c\cdot H_s)$ of glued solutions for $\phi\circ\psi$ for $c\cdot H_s$. The count of all $\partial\mathcal{M}$ proves the existence of the chain homotopy mentioned in Figure 4 like in the proof of Theorem 89. We need to justify why we can ignore those $\mathcal{M}$ which contain a solution which escapes M'. This is proved like Lemma 77, since the glued solutions for $c\cdot H_s$ either lie in M or escape M', by combining Lemmas 73, 74. So $\phi\circ\psi$ is chain homotopic to the map we get at $c=0$, which is the identity since it counts isolated Morse continuation solutions for a constant homotopy (14.1). Similarly $\psi\circ\phi\simeq \mathrm{id}$. □

14.5. **Proof of Theorem 14.** Let $H = H^\delta, H_s, f$ be as above. Let S be a model surface with weights $h_a, k_b \le 1$ (see 15.1). Denote by

$$\phi_a : SH^*(h_a H^\delta) \to MH^*(f_a) \quad\text{and}\quad \psi_b : MH^*(f_b) \to SH^*(k_b H^\delta)$$

the ϕ-,ψ-maps obtained for $h_a\cdot H_s$, $k_b\cdot H_{-s}$ and Morse functions f_a, f_b obtained by generically perturbing f. Let S' be the graph associated to S (15.19, 15.20). Then

$$\psi_S : \otimes_b SH^*(k_b H^\delta) \to \otimes_a SH^*(h_a H^\delta) \quad\text{and}\quad \psi_{S'} : \otimes_b MH^*(f_b) \to \otimes_a MH^*(f_a).$$

Lemma 79. $\otimes_a\phi_a\circ\psi_S\circ\otimes_b\psi_b$ *is chain homotopic to* $\psi_{S'}$.

Proof. The proof follows like Theorem 78 using the fact that glued solutions of $\otimes_a\phi_a\circ\psi_S\circ\otimes_b\psi_b$ either lie in M or escape M', combining Lemmas 81, 73, 74. □

Similarly, $\otimes_a\psi_a\circ\psi_{S'}\circ\otimes_b\phi_b$ is chain homotopic to ψ_S, in the obvious notation. Theorem 14 then follows upon taking direct limits and using Theorems 93, 78.

15. APPENDIX: TQFT STRUCTURE ON $SH^*(M)$ AND $HW^*(L)$

15.1. **Model Riemann surfaces.** See Figure 6. Let (S,j) be a Riemann surface with $p\ge 1$ negative punctures and $q\ge 0$ positive punctures, with a fixed choice of complex structure j and a fixed choice of parametrization $(-\infty,0]\times S^1$ and $[0,\infty)\times S^1$ respectively near the negative and positive punctures so that $j\partial_s = \partial_t$. We call these parametrizations the *cylindrical ends*. We assume S has no boundary and that away from the ends S is compact.



FIGURE 6. (S,j), genus $g=2$, $p=3$ negative and $q=2$ positive punctures, cylindrical parametrizations near the puncture (ends).

We also fix a one-form β on S satisfying $d\beta\le 0$, such that on each cylindrical end β is some positive constant multiple of dt. These constants $h_1,\ldots,h_p,k_1,\ldots k_q > 0$ will be called *weights*. The reason for using β is that we do not have a global form

dt for a general surface S, unless S is a cylinder. The condition $d\beta \leq 0$ is crucial to prove a maximum principle and an priori energy estimate (sections 15.3 and 15.4).

By Stokes's theorem, $0 \geq \int_S d\beta = \sum k_b - \sum h_a$. This is the only obstruction to finding a one-form β interpolating between the $h_1 dt, \ldots, k_q dt$ used at the ends. This explains why $p = 0$ is not allowed. Also, there is a contractible set of choices of complex structures j on S extending the j on the ends ($j\partial_s = \partial_t$), see [32, 2.2.1].

15.2. **Floer solutions.** These are smooth maps $u : S \to \overline{M}$ converging to Hamiltonian orbits as $s \to \pm\infty$ on the ends, with $du - X \otimes \beta$ holomorphic: $(du - X \otimes \beta)^{0,1} = 0$. On a cylindrical end with weight c this is Floer's equation $\partial_s u + J(\partial_t u - cX) = 0$ for the Hamiltonian cH. So u converges to 1-periodic orbits of $h_a H, k_b H$.

Let $\mathcal{M}(x_1, \ldots, x_p; y_1, \ldots, y_q; S, \beta)$ denote the moduli space of Floer solutions which converge to Hamiltonian orbits x_a at the negative ends and y_b at the positive ends. We abbreviate it by $\mathcal{M}(x_a; y_b; S, \beta)$ and call x_a, y_b the *asymptotics*. We write $\mathcal{M}_k(x_a; y_b; S, \beta)$ for the k-dimensional part of $\mathcal{M}(x_a; y_b; S, \beta)$.

Example 80. *Let S be the cylinder $Z = \mathbb{R} \times S^1$ with $p = q = 1$, weights $m' \geq m$, $\beta = f(s)\, dt$ with $f'(s) \leq 0$. We call these* continuation cylinders*. Then $u \in \mathcal{M}(x; y; Z, \beta)$ solves $\partial_s u + J(\partial_t u - f(s) X) = 0$, so it is a Floer continuation solution for the homotopy $H_s = f(s)\, H$ from $m'H$ to mH. The monotonicity condition $\partial_s h_s' = f'(s)\, h' \leq 0$ of 2.2 is equivalent to the condition $d\beta = f'(s)\, ds \wedge dt \leq 0$ of 15.1. Thus the count of $\mathcal{M}_0(x; y; Z, \beta)$ is the continuation map $SH^*(mH) \to SH^*(m'H)$.*

15.3. **Energy of Floer solutions.** Define the energy of a map $u : S \to \overline{M}$ by

$$E(u) = \frac{1}{2} \int_S \|du - X \otimes \beta\|^2 \operatorname{vol}_S.$$

Observe that for $\beta = dt$ on a cylinder this reduces to $\frac{1}{2} \int \|\partial_s u\|^2 + \|\partial_t u - X\|^2$, which equals the usual energy $\int |\partial_s u|^2 \, ds \wedge dt$ when u is a Floer solution.

For $Y \in \operatorname{Hom}(TS, u^*TM)$, the equation $Y^{0,1} = 0$ means $Y \circ j = J \circ Y$, and so $\|Y\|^2 = \omega(Y\partial_s, JY\partial_s) + \omega(Y\partial_t, JY\partial_t) = 2\omega(Y\partial_s, Y\partial_t)$ where $s + it$ is a local holomorphic coordinate on S, with $j\partial_s = \partial_t$. Thus, writing $\beta = \beta_s ds + \beta_t dt$ in components, the Floer equation $(du - X \otimes \beta)^{0,1} = 0$ implies

$$\begin{aligned} \tfrac{1}{2}\|du - X \otimes \beta\|^2 &= \omega(\partial_s u - X\beta_s, \partial_t u - X\beta_t) \\ &= \omega(\partial_s u, \partial_t u) + dH(\partial_t u)\beta_s - dH(\partial_s u)\beta_t \\ &= \frac{u^*\omega - u^*dH \wedge \beta}{ds \wedge dt}. \end{aligned}$$

Rewriting $-u^*dH \wedge \beta = -d(u^*H\, \beta) + u^*H\, d\beta$ and using the assumptions $H \geq 0$, $d\beta \leq 0$, by Stokes's theorem we obtain an a priori energy bound for Floer solutions

$$\begin{aligned} E(u) &= \textstyle\int_S u^*\omega - u^*dH \wedge \beta \\ &\leq \textstyle\int_S u^*\omega - d(u^*H\, \beta) \\ &= \textstyle\int_S d(u^*\theta - u^*H\, \beta) \\ &= \sum_{\text{negative ends } x_a} A_{h_a H}(x_a) - \sum_{\text{positive ends } y_b} A_{k_b H}(y_b). \end{aligned}$$

15.4. **Maximum principle.** In a local holomorphic coordinate $s + it$ on S, with $j\partial_s = \partial_t$ and $\beta = \beta_s ds + \beta_t dt$, the equation $(du - X \otimes \beta)^{0,1} = 0$ translates to $(du - X \otimes \beta) \circ j = J \circ (du - X \otimes \beta)$, and evaluating at ∂_s, ∂_t yields

$$\begin{aligned} \partial_t u &= X\beta_t + J\partial_s u - JX\beta_s \\ \partial_s u &= X\beta_s - J\partial_t u + JX\beta_t. \end{aligned}$$

Lemma 81. *If on the collar $H = h(R)$ and J is of contact type, then for any local Floer solution u landing in the collar, the coordinate $R \circ u$ cannot have local maxima unless it is constant. In particular, Floer solutions with asymptotics x_k are entirely contained in the region $R \leq \max R(x_k)$. The result holds even if H is time-dependent near the ends (where β is a multiple of dt). The same holds if $H = h_z(R)$ depends on $z = s + it$ on the ends, provided it is monotone: $\partial_s h'_z \leq 0$.*

Proof. Let ρ denote the R-component of $u(s,t)$. Using $\theta = -dR \circ J$ on the collar,

$$\begin{aligned}
\partial_s \rho &= dR(\partial_s u) = dR(X\beta_s - J\partial_t u + JX\beta_t) \\
&= \theta(\partial_t u) - \theta(X)\beta_t \\
\partial_t \rho &= dR(\partial_t u) = dR(X\beta_t + J\partial_s u - JX\beta_s) \\
&= -\theta(\partial_s u) + \theta(X)\beta_s \\
d^c \rho &= d\rho \circ j = (\partial_t \rho)\, ds - (\partial_s \rho)\, dt \\
&= -\theta(\partial_s u)\, ds - \theta(\partial_t u)\, dt + \theta(X)\beta_s ds + \theta(X)\beta_t dt \\
&= -u^*\theta + \theta(X)\beta \\
-dd^c \rho &= u^*\omega - d(\theta(X)\beta) \\
&= \tfrac{1}{2}\|du - X \otimes \beta\|^2\, ds \wedge dt + u^* dH \wedge \beta - d(\theta(X)\beta).
\end{aligned}$$

On the collar $X = h'\mathcal{R}$, where $\mathcal{R}$ is the Reeb vector field, so $\theta(X) = Rh'$. Thus,

$$u^* dH \wedge \beta - d(\theta(X)\beta) = h' d\rho \wedge \beta - d(\rho h' \beta) = -\rho h'' d\rho \wedge \beta - \rho h' d\beta.$$

Thus $\Delta \rho\, ds \wedge dt = -dd^c \rho \geq -\rho h'' d\rho \wedge \beta - \rho h' d\beta$, so

$$\Delta \rho + \text{first order terms} \geq -\frac{\rho h' d\beta}{ds \wedge dt}$$

Since $d\beta \leq 0$, $h' \geq 0$, $\rho \geq 0$, the right side is ≥ 0. The claim follows by the maximum principle [15] for the elliptic operator $L = \Delta + Rh''(R)\beta_t \partial_s - Rh''(R)\beta_s \partial_t$ since $Lu \geq 0$. If $H = h_t$ on an end where $\beta = C\, dt$ ($C > 0$), the extra term $-\rho(\partial_t h'_t) dt \wedge \beta = 0$. If $H = h_z$ the extra term $-\rho(\partial_s h'_z)C \geq 0$ provided $\partial_s h'_z \leq 0$. □

15.5. **Fredholm theory.** For all $u \in \mathcal{M}(x_a; y_b; S, \beta)$, Lemma 81 proves that u lies in the compact $R \leq \max(R(x_a), R(y_b))$ in $\overline{M}$. So the following Lemmas don't notice that $\overline{M}$ is non-compact, so the proofs known for closed manifolds apply. After a small generic time-dependent perturbation of (H, J) on the ends, the following hold:

Lemma 82. [32, 2.5.7]. *Solutions of $(du - X \otimes \beta)^{0,1} = 0$ are smooth and those of finite energy converge exponentially fast to Hamiltonian orbits at the ends, that is near each end $|\partial_s u| \leq ce^{-\delta|s|}$ for some constants $c, \delta > 0$.*

Lemma 83. [32, 3.1.31, 3.3.11] *or* [6, Prop.4]. *Let u be a Floer solution. Let D_u be the linearization at u of the operator defining $(du - X \otimes \beta)^{0,1} = 0$ (see 15.7), then D_u is Fredholm with* $\operatorname{index} D_u = \sum |x_a| - \sum |y_b| + 2n(1 - g - p)$, *where g is the genus of S (after filling in the punctures).*

15.6. **Smoothness of the Moduli Spaces.** For closed manifolds M, $d\beta \leq 0$ was not necessary. So the natural approach [32] was to pick any Hamiltonians H_a, H_b on the ends and any tensor $\kappa = X \otimes \beta$ which equals X_{H_a}, X_{H_b} on the ends with $\kappa = 0$ away from the ends (making u J-holomorphic there: $du^{1,0} = 0$). Schwarz [32, 4.2.17, 4.2.20] proved smoothness of $\mathcal{M}(x_a; y_b; S, \beta)$ for generic κ close to 0 away from the ends. For Liouville domains M this cannot be done, since the a priori energy estimate fails without $d\beta \leq 0$. So we need to reprove the smoothness.

Just like for Floer continuation solutions, we must allow J to depend on $z \in S$ away from the ends. So $J \in C^\infty(S \times \overline{M}, T\overline{M})$, and for each $z \in S$, $J(z,\cdot)$ is of contact type at infinity. So given H, J as in 2.3 and 1.1, we first make a small time-dependent perturbation of (H,J) on the ends to make the asymptotic Hamiltonian orbits isolated (and non-degenerate). Then in 15.7 we make a small z-dependent perturbation away from the ends, to make J *regular* (which we now explain).

The equation $(du - X \otimes \beta)^{0,1} = 0$ corresponds to $\overline{\partial} u = 0$ for the operator

$$\overline{\partial} = \tfrac{1}{2}\{(du - X \otimes \beta) + J \circ (du - X \otimes \beta) \circ j\}.$$

So the moduli space $\mathcal{M}(x_a; y_b; S, \beta)$ is the intersection of the section $\overline{\partial}$ and the zero section of a suitable Banach bundle (for details on this bundle see [32, 2.2.4]):

$$\mathcal{M}(x_a; y_b; S, \beta) = \overline{\partial}^{\,-1}(0).$$

Regularity of J means that $\overline{\partial}$ is transverse to the zero section and so $\mathcal{M}(x_a; y_b; S, \beta)$ is a smooth manifold by the Sard-Smale theorem. Regularity of J is equivalent to the surjectivity of the linearizations D_u of $\overline{\partial}$ at Floer solutions u, in particular the index of D_u will then be the dimension of the kernel of D_u. This kernel is the tangent space of $\overline{\partial}^{\,-1}(0)$, so by Lemma 83:

$$\dim \mathcal{M}(x_a; y_b; S, \beta) = \sum |x_a| - \sum |y_b| + 2n(1 - g - p).$$

15.7. **Existence of regular** J**.** We claim that for generic (in the sense of Baire) J, *which depend on* S, regularity holds: the linearization D_u of $\overline{\partial}$ is surjective.

Let (S, j, β, J) be the data described in 15.1. Pick H linear at infinity (see 17). Make a time-dependent perturbation of (H, J) on the ends to ensure that the Hamiltonian orbits that can arise as asymptotics are isolated (and non-degenerate).

By Lemma 81 all Floer solutions lie in a compact $K \subset \overline{M}$ determined by the asymptotics. We will achieve transversality by perturbing J on K, keeping J of contact type outside of K (so the maximum principle still holds outside K).

Let $D_u : W^{1,\mathfrak{p}}(u^*TM) \to L^{\mathfrak{p}}(\mathrm{Hom}^{0,1}(TS, u^*TM))$ denote the linearization of $\overline{\partial}$ at $u \in \mathcal{M}(x_a; y_b; S, \beta)$ ($\mathfrak{p} > 2$). We claim that after a generic perturbation of J, all D_u are surjective (equivalently: the D_u are transverse to the zero section).

Consider $\overline{\partial}$ for all choices of J simultaneously, where J can depend on $z \in S$. Let $\mathcal{J}$ denote the collection of all J, appropriately topologized as a Banach bundle: we require that J_z converges to the given J on the ends with superexponential convergence in any C^k-topology ([32, 4.2.11]). Linearizing at a solution $\overline{\partial}_J(u) = 0$:

$$\begin{gathered} F : T_J\mathcal{J} \oplus W^{1,\mathfrak{p}}(u^*TM) \longrightarrow L^{\mathfrak{p}}(\mathrm{Hom}^{0,1}(TS, u^*TM)), \\ F(Y, \xi) = \tfrac{1}{2}\, Y_{z,u(z)} \circ (du - X \otimes \beta)_{z,u(z)} \circ j_z + D_u \xi, \end{gathered}$$

where $T_J\mathcal{J}$ is the tangent space (infinitesimal perturbations of J with superexponential decay on the ends).

Lemma 84. *If F is surjective, there is a Baire second category subset $\mathcal{J}_{reg}(S, \beta) \subset \mathcal{J}$ of regular J (those for which D_u is surjective for all $u \in \mathcal{M}(x_a; y_b; S, \beta, J)$).*

Sketch proof. This is a standard trick, e.g. see [24, 3.1.5, p.51]. One considers the universal moduli space $\mathcal{M}$ of all Floer solutions $u \in \mathcal{M}(x_a; y_b; S, \beta, J)$ for all $J \in \mathcal{J}$. Let $\pi : \mathcal{M} \to \mathcal{J}$ denote the projection. One checks that the kernel and cokernel of $d\pi$ are isomorphic to those of D_u. Thus $d\pi$ is Fredholm, and it is onto whenever D_u is onto. Thus the regular values of π correspond to the regular J. By the Sard-Smale theorem the regular values of π in $\mathcal{J}$ are of Baire second category. $\square$

Lemma 85. *If $du - X \otimes \beta \neq 0$ on a non-empty open $\subset S$, then F is surjective.*

Proof. This is a standard argument [32, 4.2.18]. By Lemma 83, D_u is Fredholm so the image has finite codimension. Since $\operatorname{Im} D_u \subset \operatorname{Im} F$, the same is true for the image of F. Thus it suffices to show that the orthogonal complement to the image $\operatorname{Im} F$ is zero. Suppose by contradiction there is an $\eta \neq 0$ in the dual space, $L^{\mathfrak{q}}(\mathrm{Hom}^{0,1}(TS, u^*TM))$, which is perpendicular to $\operatorname{Im} F$.

Then η it is perpendicular to $\operatorname{Im} D_u$: $\langle D_u \xi, \eta \rangle_{L^2} = 0$ for all ξ. So η is a weak solution of $D_u^* \eta = 0$, and by elliptic regularity [32, 2.5.3] η is smooth. By the Carleman similarity principle [18, Cor 2.3] η can vanish at most on a discrete set of points in S. Since η is perpendicular to $\operatorname{Im} F$ and $\langle D_u \xi, \eta \rangle_{L^2} = 0$,

$$0 = \langle F(Y, \xi), \eta \rangle_{L^2} = \tfrac{1}{2} \langle Y(du - X \otimes \beta) \circ j, \eta \rangle_{L^2} \text{ for all } Y, \xi.$$

Pick $z_0 \in S$ in the open subset where $du - X \otimes \beta \neq 0$ such that $\eta_{z_0} \neq 0$. Then one can pick [31, 8.1] a vector Y at $(z_0, u(z_0))$ such that $g(Y(du - X \otimes \beta) j, \eta) > 0$. Extend Y locally and multiply it by a cut-off function which depends on $z \in S$ to make Y globally defined so that $g(Y(du - X \otimes \beta) j, \eta) \geq 0$ everywhere and > 0 near z_0. This contradicts the vanishing of the above L^2 inner product. □

Lemma 86. *Combining the previous two Lemmas, we deduce that transversality holds unless $du - X \otimes \beta = 0$ on all of S (since being non-zero is an open condition). By making a generic time-dependent perturbation of H at each end of S, we can ensure that $du - X \otimes \beta = 0$ is impossible for all Floer solutions u.*

Proof. After a generic time-dependent perturbation of H at each end of S, the asymptotics of u can never be constant orbits (since $\mathrm{Crit}(H_t)$ will change in time, generically). Suppose $du - X \otimes \beta \equiv 0$. Then $du(TS) = X \otimes \beta(TS) \subset \mathrm{span}(X)$, so the image of u lands inside a Hamiltonian orbit v of H. When S is a disc ($p = 1$, $q = 0$), $u(S) \subset v(S^1)$ means that v is contractible within $v(S^1)$ and hence constant, which we ruled out. In the general case, since we choose the time-dependent perturbations to be different on each end (of type $H = h_t(R)$ on the collar), we can ensure that v is not a common orbit of the perturbed Hamiltonians. □

Remark 87. *If the h_a, k_b of S are $\mathbb{Q}$-independent, we can make the same perturbation of H at each end (in the proof: v has periods h_a, k_b so it would be constant).*

15.8. **Compactness of the Moduli Spaces.** We claim that the $\mathcal{M}(x_a; y_b; S, \beta)$ have compactifications whose boundaries consist of broken Floer solutions (Figures 7, 8). Since the complex surface (S, j) is fixed, away from the ends we have C^∞-local hence uniform convergence to a solution, thus breaking of Floer solutions can only occur at the cylindrical ends. On those ends, the equation turns into Floer's equation so the breaking is analytically identical to breaking of Floer trajectories.

Details of the construction of compactifications for these moduli spaces can be found in Schwarz [32, 4.3.21, 4.4.1]. His work for *closed* manifolds M applies here because: 1) all $u \in \mathcal{M}(x_a; y_b; S, \beta)$ lie in the compact $R \leq \max(R(x_a), R(y_b))$ in $\overline{M}$, by 15.4; 2) they satisfy an a priori energy estimate by 15.3; and 3) no bubbling off of J-holomorphic spheres occurs since $\omega = d\theta$ is exact.

Lemma 88. *Suppose $u_n \in \mathcal{M}_1(x_a; y_b; S, \beta)$ has no C^∞-convergent subsequence. Then, passing to a subsequence, u_n converges C^∞-locally to an isolated Floer solution u defined on the same (S, β) but with one of the asymptotes x_a, y_b changed*



FIGURE 7. Floer solutions $u_n \in \mathcal{M}(x_1, x_2, x_3; y_1, y_2; \beta)$ converging to a broken solution $v\#u \in \mathcal{M}(x_2, z) \times \mathcal{M}(x_1, z, x_3; y_1, y_2; \beta)$.

at an end. On that end, there are reparametrizations $u_n(\cdot + s_n, \cdot) \to v$ *converging* C^∞*-locally to an isolated Floer trajectory as* $s_n \to \pm\infty$ (*using the sign of the end*).

Gluing theorem: given such isolated solutions v, u, *for large* $\lambda \in \mathbb{R}$ *there is a smooth* 1*-parameter family* $u_\lambda \in \mathcal{M}_1(x_a; y_b; S, \beta)$ *converging to the broken trajectory* $v\#u$ *or* $u\#v$ *in the above sense* (*resp. for breakings at negative or positive ends*).

The Fredholm index of solutions is additive with respect to gluing, so the local dimension of moduli spaces is additive (for Floer trajectories count dim $\widehat{\mathcal{M}}$, see 1.3).

15.9. **Parametrized moduli spaces.** Given two sets of regular data (S, j_0, β_0, J_0) and (S, j_1, β_1, J_1) agreeing at the ends, let $(S, j_\lambda, \beta_\lambda, J_\lambda)$ interpolate between them (while keeping the same data j, β, J near the ends). This is always possible since the choices of j, β, J each form a contractible set. Again one proves that for a generic choice of the interpolation J_λ the relevant operator $\overline{\partial}$ is transverse to the zero section and thus the moduli space $\cup_\lambda \mathcal{M}(x_a; y_b; S, \beta_\lambda, j_\lambda, J_\lambda)$ is smooth. The compactness results of section 15.8 also carry over to this setup [32, 5.2.3]. Indeed near the ends we are not varying j, β, J so the same breaking of solutions applies, and away from the ends the energy estimate forces C^∞-local hence uniform convergence.

15.10. **TQFT operations.** Fix (S, j, β) and an $H \in C^\infty(\overline{M}, \mathbb{R})$ as in 2.3. Define

$$\psi_S : SC^*(k_1 H) \otimes \cdots \otimes SC^*(k_q H) \to SC^*(h_1 H) \otimes \cdots \otimes SC^*(h_p H)$$

on the generators by counting isolated Floer solutions

$$\psi_S(y_1 \otimes \cdots \otimes y_q) = \sum_{u \in \mathcal{M}_0(x_1, \ldots, x_p; y_1, \ldots, y_q; S, \beta)} \epsilon_u \, x_1 \otimes \cdots \otimes x_p,$$

where $\epsilon_u \in \{\pm 1\}$ are orientation signs (Appendix 16). Then extend ψ_S linearly. When gradings are defined, degree$(\psi_S) = \sum |x_a| - \sum |y_b| = -2n(1 - g - p)$ by 15.6.

Theorem 89. *The* ψ_S *are chain maps, and over a field (15.17) they yield maps* $\psi_S : \otimes_{b=1}^{q} SH^*(k_b H) \to \otimes_{a=1}^{p} SH^*(h_a H)$ *which are independent of the choice of the data* (β, j, J) *relative to the ends.*

Proof. To prove that ψ_S is a chain map consider the 1-dimensional $\mathcal{M}_1(x_a; y_b; S, \beta)$. A sequence of Floer solutions approaching the boundary will break giving rise to an isolated Floer trajectory at one of the ends (15.8, Figure 7). This is precisely the definition of the Floer differential ∂ on the tensor products (16.15). Since the oriented count of the boundary components of a compact 1-dimensional manifold is zero, and after checking orientation signs in 16.15, we conclude that $\psi_S \circ \partial = \partial \circ \psi_S$. Thus ψ_S descends to the cohomology, and by the Künneth theorem we obtain

$$\psi_S : \otimes_b SH^*(k_b H) \to \otimes_a SH^*(h_a H).$$

To prove independence of the auxiliary data (β, j, J) consider the 1-dimensional part of $\bigcup_{0\leq\lambda\leq 1}\mathcal{M}(x_a; y_b; S_\lambda)$ as in 15.9, where $S_\lambda = (S, \beta_\lambda, j_\lambda, J_\lambda)_{0\leq\lambda\leq 1}$ is a regular homotopy of the data (β, j, J) relative ends. We show that ψ_{S_0} and ψ_{S_1} are chain homotopic, so the claim follows. We run the usual invariance proof of Floer homology [29]. The boundaries of the moduli spaces are of two types: either $\lambda \to 0$ or 1, which respectively yield contributions to $-\psi_{S_0}$ and ψ_{S_1}, or $\lambda \to \lambda_0 \in (0, 1)$. In the latter case the boundary is a broken solution consisting of a Floer trajectory and a "rogue" solution in $\mathcal{M}_{-1}(x_a; y_b; S_\lambda)$. Generically $\mathcal{M}_{-1}(x_a; y_b; S_\lambda)$ is empty since it has virtual dimension -1, however at finitely many λ there may exist a "rogue" solution. Let K denote the oriented count of these rogue solutions. Thus these broken solutions contribute to $K \circ \partial_0$ or $\partial_1 \circ K$ depending on the order of the breaking. The oriented count of the boundaries of a compact 1-manifold is zero, so $-\psi_{S_0} + \psi_{S_1} + K \circ \partial_0 + \partial_1 \circ K = 0$. So K is the required chain homotopy. □

15.11. **Gluing surfaces yields compositions of operations.** Two surfaces S, S' as in 15.1 can be glued along opposite ends carrying equal weights. Pick a gluing parameter $\lambda \gg 0$. Suppose a positive end $[0,\infty)\times S^1$ of S and a negative end $(-\infty, 0]\times S^1$ of S' carry the same weight. Cut off $(\lambda,\infty)\times S^1$ from S and $(-\infty,-\lambda)\times S^1$ from S'. Then glue their boundaries $[0,\lambda]\times S^1 \cup [-\lambda, 0]\times S^1$ via $(\lambda, t)\sim(-\lambda, t)$. As the weights agree, for $\lambda \gg 0$ we get a 1-form β_λ on the glued surface S_λ from the β's on the S, S' parts of S_λ. The result is a 1-family of surfaces $S_\lambda = S\#_\lambda S'$. We can glue several pairs of ends and we can also glue a positive end of S to one of its own negative ends (provided the glued surface has a negative puncture).

Lemma 90. *Let* $\mathcal{M} \subset \mathcal{M}_1(x_a; y_b; S, \beta)$ *be a connected component. Near* $\partial\mathcal{M}$*,* $\mathcal{M}$ *is parametrized by* u_λ *defined on a gluing* $S_\lambda = S\#_\lambda(\textit{infinite cylinder}, \beta = C\,dt)$. *As* $\lambda \to \infty$*,* u_λ *converges to a broken solution, broken at the gluing end.*


$\mathcal{M}$
u
u_λ
$u_\lambda(\lambda, \cdot)$
$u_\lambda(\cdot + 2\lambda, \cdot)$
v

FIGURE 8. Floer solutions u_λ converging to a broken solution $u\#v$.

Proof. Observe Figure 8. We will deal with the case of a breaking $u\#v$ on a positive end as $\lambda \to \infty$, the other case is analogous. By Lemma 88, the Floer solutions in $\mathcal{M}$ near $u\#v$ are parametrized by a gluing parameter $\lambda \gg 0$. Indeed, by an implicit function theorem argument, they can be described by a gluing construction which "glues" $u(s,t)|_{s\leq\lambda}$ with $v(s - 2\lambda, t)|_{s\geq\lambda}$. In particular, the glued solution u_λ will converge C^∞-locally to u, and on the breaking end the rescaled map $u_\lambda(s+2\lambda, t)$ will converge C^∞-locally to $v(s,t)$. The Hamiltonian orbit $v(-\infty, t)$ where the breaking occurs is the limit of $u_\lambda(\lambda, t)$ (dotted line in in Figure 8). Let $S_\lambda = S\#_\lambda(\mathbb{R}\times S^1)$ glued at the end where the breaking occurs, extending $\beta = C\,dt$ to the cylinder. "Extend" u_λ to S_λ so that it equals u_λ on the S-part and it equals $u_\lambda(s + 2\lambda, t)$ at $(s,t) \in [-\lambda,\infty)\times S^1$. This is well-defined since we identified $(\lambda, t) \sim (-\lambda, t)$. □

Theorem 91. *Let $S_\lambda = S\#_\lambda S'$ with all positive ends of S glued to all negative ends of S' in the given order. Then on the chain level $\psi_{S_\lambda} = \psi_S \circ \psi_{S'}$ for $\lambda \gg 0$.*

Remark 92. *If we do not glue all ends, the composite $\psi_S \circ \psi_{S'}$ is ill-defined. The remedy is to take disjoint unions with cylinders $Z = (\mathbb{R} \times S^1, \beta = dt)$ (which induce the identity map by Lemma 2) so that the resulting surfaces can be fully glued.*

Proof. $\psi_S \circ \psi_{S'}$ counts isolated broken Floer solutions $u\#u'$ which broke at the ends where S, S' get glued. Arguing as in Lemmas 88 and 90, there is a 1-parameter family u_λ of Floer solutions on S_λ parametrizing the Floer solutions close to $u\#u'$.

Conversely, as $\lambda \to \infty$ a 1-parameter family of isolated Floer solutions u_λ on S_λ must converge to a broken solution counted by $\psi_S \circ \psi_{S'}$. Indeed, the u_λ have a priori bounded energy depending on the fixed asymptotics, so by Lemma 82 they must break at the stretched ends. Since the u_λ are isolated, for dimensional reasons they cannot break at other ends and they only break once on each stretched end.

Since the Hamiltonian is linear at infinity, the relevant chain complexes are finitely generated and the union of the moduli spaces defining the operations ψ_{S_λ}, ψ_S, $\psi_{S'}$ is finite. So for large λ there is a bijection between the moduli spaces counted by ψ_{S_λ} and those counted by $\psi_S \circ \psi_{S'}$. We prove in 16.14 that the solutions are counted with the same orientation signs. Thus $\psi_{S_\lambda} = \psi_S \circ \psi_{S'}$ for large λ. □

15.12. TQFT operations on $SH^*(M)$.

Theorem 93. *There is a commutative diagram,*

$$\begin{tikzcd}
SH^*(k_1H) \otimes \cdots \otimes SH^*(k_qH) \arrow[r, "\psi_S"] \arrow[d, "\varphi"] & SH^*(h_1H) \otimes \cdots \otimes SH^*(h_pH) \arrow[d, "\varphi'"] \\
SH^*(k_1'H) \otimes \cdots \otimes SH^*(k_q'H) \arrow[r, "\psi_{S'}"] & SH^*(h_1'H) \otimes \cdots \otimes SH^*(h_p'H)
\end{tikzcd}$$

where the vertical maps are tensors of continuation maps, and S' is the same Riemann surface as S but with a modified form β, where $h_a \le h_a'$, $k_b \le k_b'$.

Thus the operations ψ_S induce maps on the direct limits: $\psi_S : SH^*(M)^{\otimes q} \to SH^*(M)^{\otimes p}$, and these only depend on p, q and the genus of S (for S connected).

Proof. $\varphi = \psi_{\cup_b Z_b}, \varphi' = \psi_{\cup_a Z_a}$, where $\cup_b Z_b, \cup_a Z_a$ are disjoint unions of continuation cylinders. By Theorems 89 and 91, $\varphi' \circ \psi_S$ and $\psi_{S'} \circ \varphi$ both equal $\varphi_{S''}$, where S'' is the gluing of one of the disjoint unions of cylinders onto S. □

15.13. **TQFT is compatible with filtrations.** We can filter $SC^*(H) = SC^*_\gamma(H)$ by the homology classes $\gamma \in H_1(\overline{M})$ of the generators. The Floer differential preserves the filtration, and so do Floer operations on a cylinder and a cap. The pair of pants product is additive: $SH^i_\gamma(M) \otimes SH^j_{\gamma'}(M) \to SH^{i+j}_{\gamma+\gamma'}(M)$, and similarly for the case $p = 2, q = 1$. For general S, decompose S as in 15.18. Thus:

$$\psi_S : \otimes_b SH^*_{\gamma_b}(M) \to \otimes_a SH^*_{\gamma_a}(M) \text{ is zero unless } \textstyle\sum \gamma_a = \sum \gamma_b.$$

We can also filter $SH^*(M)$ by the free homotopy classes $\gamma \in [S^1, M]$ of the generators. Then the above holds if we replace sums $\sum$ by concatenation of free loops.

In particular it is possible to restrict to the component of contractible loops: $\psi_S : SH^*_0(M)^{\otimes q} \to SH^*_0(M)^{\otimes p}$ and $c^* : H^*(M) \to SH^*_0(M) \subset SH^*(M)$.

15.14. Invariance of the TQFT structure on $SH^*(M)$.

Theorem 94. *For $\varphi : \overline{M} \cong \overline{N}$ of contact type, $\varphi_*^{\otimes p} \circ \psi_{S,M} = \psi_{S,N} \circ \varphi_*^{\otimes q}$, where $\psi_{S,M}$ is the ψ_S operation on $SH^*(M)$. So φ_* respects the TQFT.*

Proof. This is proved like Lemma 3 in 2.5: pull back the data (H, θ_N, J_N) via φ^* to M and check that $SH^*(M) \cong SH^*_f(M)$ preserves the TQFT. Since $SH^*(M) \to SH^*_f(M)$ is constructed as a direct limit of continuation maps which homotope f to 0, we concatenate TQFT operations with these continuation cylinders to obtain a commutative diagram as in Theorem 93 and then take direct limits. $\square$

15.15. **The product.** The pair of pants surface P (Figure 1) defines the product

$$\psi_P : SH^i(M) \otimes SH^j(M) \to SH^{i+j}(M),\ x \cdot y = \psi_P(x, y),$$

which is graded-commutative and associative. Commutativity is a consequence of orientation signs by Lemma 103 (we only use the $\mathbb{Z}/2\mathbb{Z}$-grading on SH^*, 16.15). Associativity is proved by gluing: glue onto one positive end of P the negative end of another copy of P. The result is a genus 0 surface with $p = 1$, $q = 3$, independently of which positive end we chose, so associativity follows from Theorems 89 and 91.

15.16. **The unit.** Let $S = \mathbb{C}$, $p = 1$, $q = 0$. The end is parametrized by $(-\infty, 0] \times S^1$, $s + it \mapsto e^{-2\pi(s+it)}$. On this end, $\beta = f(s)dt$ with $f'(s) \leq 0$, with $f(s) = 1$ for $s \leq 1$ and $f(s) = 0$ for $s \approx 0$. Extend by $\beta = 0$ away from the end.
Definition. Let $e_H = \psi_C(1) \in SH^0(H)$. Define $e = \varinjlim e_H \in SH^0(M)$.


$\beta = 0$
$\beta = f(s)dt$
$\beta = dt$
J-holomorphic since $0 \cdot H = 0$
Floer's continuation equation for $f(s) \cdot H$
Floer's equation for H

Figure 9. A cap C, and its interpretation as a continuation cylinder.

Theorem 95. *e is the unit for the multiplication on $SH^*(M)$.* $\square$

Proof. By Theorems 91, 93, $\psi_P(e, \cdot) = \psi_P \circ (\psi_C(1) \otimes \text{id}) = \psi_Z = \text{id}$ (Figure 10).



Figure 10. Gluing a cap onto a pair of pants is a continuation cylinder.

Lemma 96. *e_H is a count of the isolated finite energy Floer continuation solutions $u : \mathbb{R} \times S^1 \to \overline{M}$ for the homotopy $f(s)H$ from H to 0.*

Proof. As $X_H \otimes \beta = X_{f(s)H} \otimes dt$, $(du - X \otimes \beta)^{0,1} = 0$ becomes $\partial_s u - J(\partial_t u - X_{f(s)H}) = 0$. Near $s = \infty$ such u are J-holomorphic, $\partial_s u - J\partial_t u = 0$. The finite energy condition implies that u converges to an orbit of X_H at $s = -\infty$ by Lemma 82 and that the singularity at $s = \infty$ is removable. So u extends to a J-holomorphic map over the puncture at $s = \infty$ (see [23]) recovering the solution $\mathbb{C} \to \overline{M}$ counted by ψ_C. $\square$

Lemma 97. *For $H = H^\delta$ as in Section 4, e_H = sum of the absolute minima of H.*

Proof. Pick J so that $g = \omega(\cdot, J\cdot)$ is Morse-Smale for H. Rescale H so that all Floer continuation solutions for $f(s)H$ are time-independent by Claim 5, 14.3. So e_H counts isolated Morse continuation solutions. So e_H is a linear combination of the minima of H. But e_H counts $-f(s)\nabla H$ trajectories starting from a minimium of H, so they must be constant. The claim follows. $\square$

Theorem 98. *$e = \varinjlim e_H$ is the image of* 1 *under* $c^* : H^*(M) \to SH^*(M)$.

Proof. $e_{H^\delta} = c_0^*(1)$ via $c_0^* : H^*(M) \cong MH^*(H^\delta) \equiv SH^*(H^\delta)$ by Lemma 97. Via continuation $\varphi : SH^*(H^\delta) \to SH^*(H)$, $e_H = \varphi(e_{H^\delta}) = \varphi(c_0^* 1) = c_H^*(1)$ by Theorem 93 ($c_H^* : H^*(M) \to SH^*(H)$). Now take direct limits. Claim also follows by Thm.14. □

15.17. **Unital ring structure over $\mathbb{Z}$.** So far we worked over a field $\mathbb{K}$ of coefficients (or a Novikov algebra defined over $\mathbb{K}$ in Section 6) because in 15.10 we used Künneth's theorem to pass from $H_*(\otimes_a SC^*(h_a H))$ to $\otimes_a SH^*(h_a H)$. However, operations for $p = 1$, in particular the unital ring structure, are also defined over any commutative ring since we can lift ψ_S to $\otimes_b SH^*(k_b H)$ via $\otimes_b [x_b] \mapsto \psi_S([\otimes_b x_b])$.

15.18. **A minimal set of generating surfaces under gluing.** Denote C, Z, P, Q the genus 0 surfaces with (p, q) respectively $(1, 0)$, $(1, 1)$, $(1, 2)$, $(2, 1)$. With a little care as in Figure 11 one shows that only C,Z,P,Q are needed to generate via gluing any surface with $p \geq 1$. For example, the cylinder with $(p, q) = (2, 0)$ is $Z_2 = Q\#C$ and the pair of pants $(p, q) = (3, 0)$ is $T = (Q \cup Z)\#Q\#C$. By Theorem 91:



FIGURE 11. Dark dots are punctures with the assigned sign.

Corollary 99. *Every operation* $\psi_S : SH^*(M)^{\otimes q} \to SH^*(M)^{\otimes p}$ *is a composition of operations, chosen among the four basic ones: the unit* $\psi_C : \mathbb{K} \to SH^*(M)$*, the identity* $\psi_Z = id$*, the product* ψ_P *and the coproduct* $\psi_Q : SH^*(M) \to SH^*(M)^{\otimes 2}$.

15.19. **Associated graphs.** Given a decomposition as above, associate a directed graph S' to the decomposition by replacing C, Z, P, Q by the corresponding graphs C', Z', P', Q' (Figure 12) and gluing them according to the decomposition of S.



FIGURE 12. Graphs C', Z', P', Q'. Dark dots are internal vertices. Crosses are punctures: the edge is infinite in that direction.

15.20. **Morse operations.** The Morse operations on the Morse cohomology $H^*(M)$,

$$\psi_{S'} : H^*(M)^{\otimes q} \to H^*(M)^{\otimes p},$$

are defined analogously to the TQFT operations on $SH^*(M)$, replacing S by a graph S'. Here S' is oriented and has edges parametrized by $(-\infty, 0]$, $[0, \infty)$ or $[0, \ell]$ (for finite $\ell \geq 0$) respectively called *negative ends* e_a, *positive ends* e_b and *internal edges* e_c of length ℓ. For each edge e_i pick a Morse function f_i. For critical points $x_a \in \mathrm{Crit}(f_a)$, $y_b \in \mathrm{Crit}(f_b)$ define the moduli space of *Morse solutions*, $\mathcal{M}(x_a; y_b; S', f_i)$, consisting of the continuous maps $u : S' \to \overline{M}$ which are $-\nabla f_i$ gradient flows along e_i and which converge to x_a, y_b at infinity on the edges e_a, e_b. The value of u at vertices where edges meet corresponds to points lying in the intersection of stable/unstable manifolds of the f_i. The smoothness of $\mathcal{M}(x_a; y_b; S', f_i)$ is guaranteed by making these intersections transverse by choosing the f_i generically.

These operations are well-known (Betz-Cohen [5], Fukaya [19]) and they define a TQFT structure on $H^*(M)$. The cup product on $H^*(M)$ corresponds to the Morse product operation $\psi_{P'}$, where P' is the Y-shaped graph of Figure 12 consisting of two positive edges and one negative edge all of which meet in one internal vertex.

We include the unit $1 \in H^0(M)$ in this formalism by allowing the graph C' of Figure 12 consisting of one negative edge carrying a homotopy f_s from a Morse function f to 0. Let $\psi_{C'} : \mathbb{K} \to H^*(M)$ be the count of isolated finite energy $-\nabla f_s$ flowlines along C'. The proof that $\psi_{C'}(1) = 1$ is similar to Theorem 95.

Given S as in 15.1, consider the graph S' associated to S in 15.19. The negative and positive edges are indexed by a, b respectively, and associate lengths ℓ_i to the internal edges. Pick generic Morse functions f_a, f_b on $\overline{M}$ with $-\nabla f_a, -\nabla f_b$ pointing inwards on the collar. After identifications with ordinary cohomology, we obtain

$$\psi_{S'} : H^*(M)^{\otimes q} \to H^*(M)^{\otimes p}.$$

This map does not depend on the choice of S' associated to S, or the choices ℓ_i, f_a, f_b by continuation and gluing arguments analogous to the $SH^*(M)$ case.

15.21. **Open model surfaces** S**.** Let (S, j) be a Riemann surface isomorphic to the closed unit disc with $p + q$ boundary points removed. These punctures are marked in the oriented order as $1, \ldots, p, 1, \ldots q$, the first $p \geq 1$ are called *negative*, the other $q \geq 0$ *positive*. Fix parametrizations as *strip-like ends* $(-\infty, 0] \times [0, 1]$ and $[0, \infty) \times [0, 1]$ near negative and positive punctures, such that j becomes standard: $j\partial_s = \partial_t$. Figure 13 shows two ways to think of the pair of pants $(p, q) = (1, 2)$.



Figure 13. Open-string pair of pants. Dark circles are punctures.

Fix a one-form β on S with $d\beta \leq 0$, such that on each strip-like end β is a positive constant multiple of dt. These constants $h_a, k_b > 0$ are called *weights* and are generic (not periods of Reeb chords). We require that the pullback $\beta|_{\partial S}$ vanishes ($\beta|_{\partial S}(T\partial S) = 0$) and that β is exact near ∂S.

By Stokes's theorem, $0 \geq \int_S d\beta = \sum k_b - \sum h_a$, which is the only constraint to finding β given weights. Indeed one could construct β quite canonically [3, Sec.2g].

Given two such sets of data for S, S', then as in 15.11 we can glue a positive end of S with a negative end of S' provided they carry the same weight. The resulting surface $S\#S'$ carries glued data $j\#j', \beta\#\beta'$ satisfying the above conditions.

More generally, we also allow S to be a disjoint union of surfaces as above, and we also allow surfaces S which are the result of gluing any such surfaces along several ends (provided $p \geq 1$ after gluing), so the surface can have certain holes.

15.22. **Wrapped TQFT structure.** Fix a Hamiltonian H as in the definition of $HW^*(L;H)$ with (possibly non-generic) slope 1 at infinity, and write $X = X_H$. In analogy with 15.2, define the moduli space of *wrapped solutions* $\mathcal{W}(x_a; y_b; S, \beta)$ consisting of smooth $u : S \to \overline{M}$ with $u(\partial S) \subset \overline{L}$, solving $(du - X \otimes \beta)^{0,1} = 0$, converging on the ends to the Hamiltonian chords x_a, y_b for $h_a H, k_b H$. The construction of the TQFT is analogous to the $SH^*(M)$ case, after three technical remarks:

1) Maximum principle. *All* $u \in \mathcal{W}(x_a; y_b; S, \beta)$ *have* $R(u) \leq \max(R(x_a), R(y_b))$.

Proof. The proof is identical to Lemma 81, except maxima might occur on ∂S. But then by Hopf's Lemma [15, Sec.6.4.2] the derivative $\partial_t \rho$ of $\rho = R \circ u$ in the outward normal direction ∂_t on ∂S would be positive, which it isn't:

$$\partial_t \rho = dR(\partial_t u) = dR(X\beta_t + J\partial_s u - JX\beta_s) = -\theta(\partial_s u) = 0$$

using $dR(X) = 0$, $dR \circ J = -\theta$, $\beta|_{\partial S} = 0$ so $\beta_s = 0$, and finally using $\theta|_{\overline{L}} = 0$ on the collar and $\partial_s u \in T\overline{L}$ on ∂S (since $\partial_s = -j\partial_t$ is tangent to ∂S). □

2) Energy. This is defined and bounded as in 15.3 using $\theta|_{\overline{L}} = df$ and $\beta|_{\partial S} = 0$,

$$E(u) \leq \textstyle\sum_{a=1}^{p} A_{h_a H}(x_a) - \sum_{b=1}^{q} A_{k_b H}(y_b).$$

3) Transversality. Proceed as in 15.6, so 15.7 can be applied, except we need to discuss trivial solutions $du - X \otimes \beta = 0$. Let γ be a boundary path in ∂S connecting two consecutive punctures z, z' with weights c, c'. Since the pull-back of β to ∂S vanishes and β is exact near ∂S, we can define a complex coordinate $s + it$ near γ such that: $\partial_s = \dot{\gamma}$ and $\beta = l(t)\, dt$ along γ, for some function $l(t)$ interpolating c, c'. Since $du - X \otimes \beta = 0$, $du(\dot{\gamma}) = 0$ so u is constant along γ. Thus z, z' are two (possibly equal) Hamiltonian chords such that the initial point of one is the end point of the other. This gives rise to a triple intersection in $\overline{L} \cap \varphi^c(\overline{L}) \cap \varphi^{c+c'}(\overline{L})$. As in 3.2, after perturbing $H, \overline{L}$, one can [3, Lemma 8.2] rule out these intersections provided $\dim M \geq 4$, which we tacitly assume (see [3, Sec.5b]).

These remarks imply that $\mathcal{W}(x_a; y_b; S, \beta)$ is smooth and compact up to breaking at the ends. On the ends, the equation turns into Floer's equation, so the breaking analysis is the same as for strips. Counting wrapped solutions, Theorem 93 becomes

Theorem 100. *There are TQFT maps* $\mathcal{W}_S : \otimes_b HW^*(L; k_b H) \to \otimes_a HW^*(L; h_a H)$ *which in the direct limit become TQFT operations* $\mathcal{W}_S : HW^*(L)^{\otimes q} \to HW^*(L)^{\otimes p}$, $(p \geq 1, q \geq 0)$*, which do not depend on the choices of* β, j, J.

This includes the unital ring structure: $\mathcal{W}_P : HW^*(L)^{\otimes 2} \to HW^*(L)$ with unit $\mathcal{W}_C : \mathbb{K} \to HW^*(L)$ where P is a disc with $1{+}2$ punctures and C is a disc with $1{+}0$ punctures. As in 15.15, the product is associative. It is not graded-commutative in general (the Lagrangian case lacks the symmetry of P).

15.23. $HW^*(L)$ **is a module over** $SH^*(M)$**.** Pick a surface S as in 15.21. Introduce negative and positive punctures in the interior of S. Fix cylindrical parametrizations $(-\infty, 0] \times S^1$ and $[0, \infty) \times S^1$ respectively near those punctures, with $j\partial_s = \partial_t$. Let β be a 1-form on the resulting surface S with: $d\beta \leq 0$, β is a positive multiple of dt near each puncture, β is exact near ∂S, the pull-back $\beta|_{\partial S} = 0$. Consider the moduli space of solutions $u : S \to \overline{M}$ of $(du - X \otimes \beta)^{0,1} = 0$, with $u(\partial S) \subset \overline{L}$, converging at boundary punctures to Hamiltonian chords and at interior punctures to 1-periodic Hamiltonian orbits. The maximum principle and energy estimate still hold, so transversality and compactness up to breaking on the ends are proved as



FIGURE 14. The surface D with punctures $(w'; w; x)$ viewed in two ways.

before. The count of these moduli spaces define operations involving tensor products of the groups $HW^*(L)$ and $SH^*(M)$. Consider Figure 14: at the chain level such a solution contributes $\pm w'$ to $\mathcal{W}_D(x \otimes w)$ where

$$\mathcal{W}_D : SH^*(M) \otimes HW^*(L) \to HW^*(L).$$

Thus we deduce the following result, which is known by experts:

Theorem 101. *$\mathcal{W}_D$ defines a module structure of $HW^*(L)$ over $SH^*(M)$.*

Proof. By construction $\mathcal{W}_D$ is linear in $SH^*(M)$ and $HW^*(L)$. Next we check that $\mathcal{W}_D(e, \cdot) = \mathrm{id}$ for the unit $e \in SH^*(M)$. Given H and weights, consider Figure 15: capping off the closed orbit in Figure 14 yields a strip, which we can homotope to the standard continuation strip $(\mathbb{R} \times [0,1], \beta = f(t)dt, j\partial_s = \partial_t)$ where $f'(t) \leq 0$ and f interpolates the weights. So $\mathcal{W}_D(e_H, \cdot)$ is chain homotopic to a continuation map. So, in cohomology, taking direct limits, $\mathcal{W}_D(e, \cdot)$ is the identity, as required.



FIGURE 15. Gluing a cap onto Figure 14. Gluing 2 copies of Figure 14, and finally the configuration obtained after an isotopy.

Finally we need $\mathcal{W}_D$ to be compatible with the product on $SH^*(M)$, namely $\mathcal{W}_D \circ (\mathrm{id} \circ \mathcal{W}_D) = \mathcal{W}_D \circ (\psi_P \otimes \mathrm{id})$. It suffices to check that for given H and weights, at the chain level $\mathcal{W}_D(x, \mathcal{W}_D(x', w))$ is chain homotopic to $\mathcal{W}_D(\psi_P(x, x'), w)$.

Consider the last two configurations in Figure 15: one is the gluing of two copies of Figure 14 corresponding to $\mathcal{W}_D(x, \mathcal{W}_D(x', w))$, the other is obtained by an isotopy of this one and it is the gluing corresponding to $\mathcal{W}_D(\psi_P(x, x'), w)$.

In the last figure, we first pull-back the form β via the isotopy, then we deform it (by linear interpolation) to make it equal to the β obtained from the gluing of the punctured strip with the pair-of-pants.

Now run the same argument as in the proof of Theorem 91: combining a stretching argument with the a priori energy estimate shows that each of the two configurations arises precisely from gluing broken solutions. Since the two configurations are homotopic, $\mathcal{W}_D(x, \mathcal{W}_D(x', w))$ and $\mathcal{W}_D(\psi_P(x, x'), w)$ are chain homotopic. □

16. APPENDIX 2: COHERENT ORIENTATIONS.

16.1. **Coherent orientations for Floer trajectories.** A *coherent orientation* for the $\mathcal{M}(x, y)$ of 1.3 is a continuous map associating to $u \in \mathcal{M}(x, y)$ an orientation $\sigma(u)$ of the determinant line bundle, $\mathrm{Det}\, D_u = \Lambda^{\max} \ker D_u = \Lambda^{\max} T_u \mathcal{M}(x, y)$, such that the orientations glue correctly: there is an associative gluing operation # such that $\sigma(u)\#\sigma(v) = \sigma(u\#v)$ for the broken solution $u\#v$ (which agrees with the orientation of nearby unbroken solutions under the compactification construction). Coherent orientations were first constructed by Floer-Hofer [17] as follows.

16.2. **Fredholm operators on trivial bundles over a cylinder.** One first constructs coherent orientations for families of Fredholm operators of the form

$$L : W^{1,p}(\xi) \to L^p(\xi),\ Lu = \partial_s u + J_{s,t}\partial_t u + A_{s,t}u$$

on a trivial vector bundle $E \to Z$ over the cylinder $Z = \overline{\mathbb{R}} \times S^1$ (compactified appropriately by adding two circles $S^1_\pm$ at infinity), such that $A_{s,t} \in \mathrm{End}(\xi)$ converges to self-adjoint isomorphisms $A_{\pm\infty,t}$ and $L \to L^\pm = J_{\pm\infty,t}\partial_t + A_{\pm\infty,t}$ as $s \to \pm\infty$.

Once the asymptotic operators $L^\pm$ at the ends are fixed, the family $\mathcal{O}_E(L^-, L^+)$ of such operators is a contractible (indeed convex) set [17, Prop 7]. Therefore the real line bundle $\mathrm{Det}(\mathcal{O}_E(L^-, L^+)) \to \mathcal{O}_E(L^-, L^+)$ of all determinants

$$\mathrm{Det}(L) = (\Lambda^{\max}\mathrm{coker}\, L)^\vee \otimes \Lambda^{\max} \ker L$$

is trivial, and an orientation $\sigma(L)$ is a choice of trivialization.

16.3. **Gluing the operators and coherent orientations.** For operators L, K which become constant respectively for $s \gg 0, s \ll 0$, with $L^+ = K^-$, we can glue them $L\#K$ to yield a natural isomorphism $\mathrm{Det}\, L\#K \cong \mathrm{Det}\, L \times \mathrm{Det}\, K$. Then define $\sigma(L\#K)$ by gluing two given orientations $\sigma(L)\#\sigma(K)$ via the gluing map

$$\mathcal{O}_E(L^-, L^+) \times \mathcal{O}_E(L^+, K^+) \to \mathcal{O}_E(L^-, K^+).$$

Now construct a coherent orientation as follows. Fix an operator L_0 of the type that arises at $s = -\infty$. Choose the orientation $1^\vee \otimes 1 \in \mathbb{R}^\vee \otimes \mathbb{R}$ for the isomorphism $L = \partial_s + L_0$. So we get an orientation on $\mathcal{O}_E(L_0, L_0)$. Pick any orientation of $\mathcal{O}_E(L_0, L^+)$ for all $L^+ \neq L_0$. Coherence requires the orientations on $\mathcal{O}_E(L_0, L^+), \mathcal{O}_E(L^+, L_0)$ to determine that on $\mathcal{O}_E(L_0, L_0)$, so we deduce an orientation for $\mathcal{O}_E(L^+, L_0)$. Orientations on $\mathcal{O}_E(L_0, L^-)$, $\mathcal{O}_E(L^-, L^+)$, $\mathcal{O}_E(L^+, L_0)$ determine that on $\mathcal{O}_E(L_0, L_0)$, so we deduce an orientation for $\mathcal{O}_E(L^-, L^+)$. The construction is well-defined since gluing $1^\vee \otimes 1$ with itself yields $1^\vee \otimes 1$.

16.4. **Fredholm operators on bundles $E \to \overline{M}$.** The above can be applied to any symplectic vector bundle $E \to \overline{M}$ by considering all operators L on u^*E for smooth maps $u : Z \to \overline{M}$ such that in a trivialization L is an operator of type above. The bundles u^*E have the homotopy type of S^1 so u^*E is trivial since $\pi_0(Sp(2n)) = 0$, but two choices of trivialization may be non-homotopic as $\pi_1(Sp(2n)) = \mathbb{Z}$. A careful choice [17, Lemmas 13&15] of trivializations ensures we can define an orientation for each equivalence class $[L, u]$ of operators with prescribed: asymptotic $L^\pm$, index(L), asymptotics $u(\pm\infty, t)$. As before, we construct coherent orientations $\sigma([L, u])$ for all classes $[L, u]$, with the additional features that one has to pick $\sigma(L_0, u_0)$ for every free homotopy class $u_0 \in [S^1, M]$ and one picks an orientation of $[L, u]$ for $L^\pm = L_0$ with minimal index $\mathrm{ind}(L) > 0$ (if it exists).

16.5. **Definition of orientation signs for $SH^*(M)$.** Apply 16.4 to $T\overline{M} \to \overline{M}$ to get a coherent orientation σ. Then σ is defined on all linearizations D_u of Floer's equation along a Floer trajectory u. An isolated Floer trajectory u has a natural orientation $\partial_s u$ determined by its flow. Define the $\epsilon_u \in \{\pm 1\}$ of 2.1 by $\sigma(u) = \epsilon_u \cdot \partial_s u$. For an isolated Floer continuation solution v define the $\epsilon_v \in \{\pm 1\}$ of 2.2 by $\epsilon_v = \sigma(v)$.

16.6. **Using orientation signs to prove $\partial^2 = 0$.** Let $u\#u' \in \mathcal{M}_0(x,y)\#\mathcal{M}_0(y,z)$ be a broken Floer trajectory lying at the boundary of a connected component $\mathcal{M} \subset \mathcal{M}_1(x,z)$. Floer's gluing map parametrizes a neighbourhood of this boundary by a gluing parameter $\lambda \gg 0$ as follows. It interpolates the two maps (*gluing data*)

$$(u_\lambda = u(s+\lambda,t))|_{s\leq -1} \;\text{ and }\; (u'_{-\lambda} = u'(s-\lambda,t))|_{s\geq 1}$$

to obtain an approximate solution to Floer's equation, then it finds the closest Floer solution $u\#_\lambda u' \in \mathcal{M}$ via the implicit function theorem. Recall the notation $\widehat{\mathcal{M}}$, 1.3. We can canonically orient $\widehat{\mathcal{M}}(x,y)\times\widehat{\mathcal{M}}(y,z)$ at (u,u') by the pair of basis vectors

$$((\partial_s u, -\partial_s u'),\ (\partial_s u, \partial_s u')).$$

Differentiating the gluing data $(u_\lambda, u'_{-\lambda})$ in λ and s gives rise to the two vectors $(\partial_s u_\lambda, -\partial_s u'_{-\lambda})$, $(\partial_s u_\lambda, \partial_s u'_{-\lambda})$, which agree with the above canonical orientation. Thus for large λ, the pair $(\partial_\lambda(u\#_\lambda u'), \partial_s(u\#_\lambda u'))$ defines the correct gluing orientation of $T\widehat{\mathcal{M}}(x,z)_{u\#_\lambda u'}$. Quotienting by the translations, we deduce that $\partial_\lambda(u\#_\lambda u')$ defines the glued orientation on $\mathcal{M}$ ($\cong$ interval) near the boundary. Since $u\#_\lambda u'$ approaches the boundary as λ increases, this is the outward-pointing orientation.

Thus the two boundaries $u\#u'$, $u_2\#u'_2$ of $\mathcal{M}$ are oppositely oriented by the gluing map: $\partial_s u\#\partial_s u' = -\partial_s u_2\#\partial_s u'_2$. Since they belong to the same component $\mathcal{M}$, they carry the same asymptotic operators, so $\sigma(u\#u') = \sigma(u_2\#u'_2)$. By coherence

$$\sigma(u\#u') = \sigma(u)\#\sigma(u') = \epsilon_u\epsilon_{u'}\cdot\partial_s u\#\partial_s u',$$

and similarly for $\sigma(u_2\#u'_2)$. Combining these results yields the required equality $\epsilon_u\epsilon_{u'} + \epsilon_{u_2}\epsilon_{u'_2} = 0$, which is needed to prove $d\circ d = 0$ in 2.1.

16.7. **Using orientation signs to prove $\partial\varphi = \varphi\partial$.** (where φ is a continuation map, 2.2). Let $u\#v$ and $v'\#u'$ be broken continuation solutions, where v, v' are Floer continuation solutions, u, u' are Floer trajectories. Their orientations are defined by the orientations $\partial_s u$, $\partial_s u'$ respectively. However, the gluing data is now

$$(u_{2\lambda}|_{s\leq -\lambda-1},\ v|_{s\geq -\lambda+1}) \quad\text{ and }\quad (v'|_{s\leq \lambda-1},\ u'_{-2\lambda}|_{s\geq\lambda+1}).$$

Differentiating in λ: $(2\partial_s u_{2\lambda}, 0)$, $(0, -2\partial_s u'_{-2\lambda})$. So for $\lambda\gg 0$ the glued orientations are $\partial_\lambda(u\#_\lambda v)$ (outward-pointing near the boundary) and $-\partial_\lambda(v'\#_\lambda u')$ (inward).

Suppose the two boundaries of $\mathcal{M}$ are of the same type, say $u\#v$ and $u_2\#v_2$. By coherence $\sigma(u\#v) = \sigma(u)\#\sigma(v)$, and $\sigma(u\#v) = \sigma(u_2\#v_2)$ since they lie in the same component $\mathcal{M}$. Thus, using the above gluing result, $\epsilon_u\epsilon_v = -\epsilon_{u_2}\epsilon_{v_2}$, so the two contributions to $\partial\circ\varphi$ cancel out. Similarly, if both boundaries are of type $v'\#u'$ we get two contributions to $\varphi\circ\partial$ which cancel out.

If the boundaries are $u\#v$ and $v'\#u'$, then the glued orientations either both agree or both disagree with the boundary orientation of $\mathcal{M}$. Using coherence, we deduce $\epsilon_u\epsilon_v = \epsilon_{v'}\epsilon_{u'}$, which are the required signs to prove $\partial\circ\varphi = \varphi\circ\partial$.

16.8. **Coherent orientations for Floer solutions.** For TQFT operations we need coherent orientations using smooth maps $u : S \to \overline{M}$ on a punctured Riemann surface S with prescribed parametrizations on the cylindrical ends (appropriately compactified with *asymptotic circles* S^1_a, S^1_b at the ends). We mimic the construction from symplectic field theory [6] which builds upon [14, 1.8].

16.9. **Fredholm operators over punctured surfaces.** First consider complex vector bundles $E \to S$ with prescribed trivializations on the asymptotic circles, and

$$Lu\cdot X = \nabla_X u + J\nabla_{jX}u + A_z(u)\cdot X,$$

for $L : W^{1,p}(E) \to L^p(\mathrm{Hom}^{0,1}(TS,E))$, $A \in \mathrm{Hom}(E, \mathrm{Hom}^{0,1}(TS,E))$, which on the cylindrical ends are operators of the type 16.2. Denote the space of such operators:

$$\mathcal{O}_E(L_a; L_b) = \mathcal{O}_E(L_1, \ldots, L_p; L_1, \ldots, L_q)$$

where L_a, L_b are the asymptotic operators respectively at the negative and positive ends. Just as in 16.2, $\mathcal{O}_E(L_a; L_b)$ is a contractible space so the determinant bundle over $\mathcal{O}_E(L_a; L_b)$ is trivial and an orientation is a choice of trivialization.

16.10. **Gluings and disjoint unions.** Just as in 16.3 there is a gluing operation: given $E \to S$, $E' \to S'$ with matching trivializations over the respective punctures y_c, x'_c where we glue, we obtain a glued bundle $E'' \to S''$ and a gluing map

$$\# : \mathcal{O}_E(L_a; L_b, L_{-c}) \times \mathcal{O}_{E'}(L_c, L_{a'}; L_{b'}) \to \mathcal{O}_{E''}(L_a, L_{a'}; L_b, L_{b'}),$$

where we always use the convention that L_{-c} is an abbreviation for the reversed ordering $(\ldots, L_2, L_1)$ of the operators $(L_1, L_2, \ldots)$, so inductively in c we are gluing on the c-end the pair of asymptotes $L_{-c} = L_c$. This ensures the associativity of the gluing operation $\#$ defined on the orientations $\sigma(L)$ of $\mathrm{Det}\, L$ [7, Cor 7].

The disjoint union of bundles determines a natural isomorphism

$$\mathrm{Det}\, L \otimes \mathrm{Det}\, L' \to \mathrm{Det}(L \cup L').$$

Denote $\sigma(L) \cup \sigma(L')$ the induced operation on orientations. If a gluing between operators K and $L \cup L'$ only involves gluing K with L, then *distributivity* holds:

$$\sigma(K)\#(\sigma(L) \cup \sigma(L')) = (\sigma(K)\#\sigma(L)) \cup \sigma(L').$$

16.11. **Axiomatic construction of coherent orientations.** If S has no punctures, then $\mathcal{O}_E(\emptyset; \emptyset)$ contains a Cauchy-Riemann operator $\overline{\partial}$, which is $\mathbb{C}$-linear. Thus $\ker \overline{\partial}$, $\mathrm{coker}\, \overline{\partial}$ are complex, so they are canonically oriented. Therefore, $\overline{\partial}$ defines a *canonical orientation* $\sigma(\overline{\partial})$ on $\mathcal{O}_E(\emptyset; \emptyset)$.

Now consider the standard trivial bundles triv $\to \mathbb{C}$. Denote the operators L on these bundles by $D^{\pm}$ depending on whether ∞ is a positive or negative puncture. Define an orientation on each $\mathcal{O}_{\mathrm{triv}\to\mathbb{C}}(A; \emptyset)$ by fixing an operator D^- and picking an orientation for it. This determines an orientation for any D^+ by coherence: gluing an appropriate D^- forces $\sigma(D^-)\#\sigma(D^+)$ to be the canonical orientation.

Since we established orientations for any $D_i^- \in \mathcal{O}_{\mathrm{triv}\to\mathbb{C}}(A_i; \emptyset)$, we can define

$$\sigma(D_1^- \cup \cdots \cup D_k^-) = \sigma(D_1^-) \cup \cdots \cup \sigma(D_k^-),$$

and similarly for $D_i^+ \in \mathcal{O}_{\mathrm{triv}\to\mathbb{C}}(\emptyset; A_i)$.

Definition 102. *The orientation $\sigma(K)$ for $K \in \mathcal{O}_E(L_a; L_b)$ is defined by capping off the punctures and requiring that we obtain the canonical orientation:*

$$\sigma(\cup D_{-a}^+)\#\sigma(K)\#\sigma(\cup D_{-b}^-) = \textit{canonical},$$

with the convention that D_{-a} is the reverse ordering $(\ldots, D_2, D_1)$ of the D_a.

16.12. **Orientation signs arising from gluing.** We now want to find out whether a glued orientation $\sigma(L)\#\sigma(L')$ agrees with $\sigma(L\#L')$ or not.

Lemma 103. [6, Thm.2] *If we exchange the position of two consecutive b-labels $k, k{+}1$, the coherent orientation of $\mathcal{O}_E(L_a; L_b)$ changes by $(-1)^{\mathit{ind}\, D_k \cdot \mathit{ind}\, D_{k+1}}$, where the $D_b \in \mathcal{O}_{\mathit{triv}\to\mathbb{C}}(L_b; \emptyset)$ cap off the b-ends. Similarly for a-labels using the $D_a \in \mathcal{O}_{\mathit{triv}\to\mathbb{C}}(\emptyset; L_a)$. For a general permutation of the ends, iterate this result.*

Sketch proof. For simplicity, we consider the pair of pants case. We want to compare $L \in \mathcal{O}_E(L_1; L_2, L_3)$ and $L' \in \mathcal{O}_E(L_1; L_3, L_2)$, so we cap off the ends $2, 3$ so that both operators now lie in $\mathcal{O}_{E'}(L_1;\emptyset)$ so they are equally oriented. Thus $\sigma(L\#(D_3\cup D_2)) = \sigma(L'\#(D_2\cup D_3))$. So, by our axiomatic construction,

$$\sigma(L)\#(\sigma(D_3)\cup\sigma(D_2)) = \sigma(L')\#(\sigma(D_2)\cup\sigma(D_3)).$$

Thus $\sigma(L)$, $\sigma(L')$ are the same if and only if $\sigma(D_2)\cup\sigma(D_3)$ and $\sigma(D_3)\cup\sigma(D_2)$ are the same. But for these caps, we are switching the order of the two brackets of

$$(\Lambda^{\max}\mathrm{coker}^\vee D_2 \otimes \Lambda^{\max}\ker D_2)\otimes(\Lambda^{\max}\mathrm{coker}^\vee D_3\otimes\Lambda^{\max}\ker D_3),$$

so the orientation changes by a sign: $(-1)^{(\dim\ker D_2+\dim \mathrm{coker} D_2)\cdot(\dim\ker D_3+\dim\mathrm{coker} D_3)}$ $= (-1)^{(\dim\ker D_2-\dim\mathrm{coker} D_2)\cdot(\dim\ker D_3-\dim\mathrm{coker} D_3)} = (-1)^{\mathrm{ind} D_2\cdot \mathrm{ind} D_3}$ □

Lemma 104 (Prop.8, Prop.10, [6])**.** *$\sigma(L)\#\sigma(L')$ agrees with $\sigma(L\#L')$ when all ends are glued, that is for gluings of type $\mathcal{O}_E(L_a;L_{-c})\times\mathcal{O}_{E'}(L_c;L_{b'})\to\mathcal{O}_{E''}(L_a;L_{b'})$. For partial gluings, $\mathcal{O}_E(L_a;L_b,L_{-c})\times\mathcal{O}_{E'}(L_c,L_{a'};L_{b'})\to\mathcal{O}_{E''}(L_a,L_{a'};L_b,L_{b'})$, the orientations $\sigma(L\#L')$ and $\sigma(L)\#\sigma(L')$ differ by $(-1)^{\sum \mathrm{ind}\, D_b\cdot\sum\mathrm{ind}\, D_{a'}}$. (Mnemonically: we are reordering $L_b;L_{a'}$ to $L_{a'};L_b$).*

Sketch proof. The key behind the first claim is that for complex linear operators, gluings preserve the complex orientations. Now for general L, L', we can assume the ends labelled a, b' are capped off since they don't matter. Then attach cylinders Z_{-c}, Z'_c at the c, c' ends so that the new asymptotic operators are complex linear. The resulting operators $L\cup Z_{-c}$, $Z'_c\cup L'$ are now homotopic to complex linear ones, so they glue well: $\sigma((L\cup Z_{-c})\#(Z'_c\cup L')) = \sigma(L\cup Z_{-c})\#\sigma(Z'_c\cup L')$. Finally, by the associativity of $\#$, we can move the Z'_c over to the Z_{-c} and "cancel them off" in pairs, to conclude the first claim: $\sigma(L\cup L') = \sigma(L)\#\sigma(L')$.

For the second claim, write $\sigma_a = \cup_a\sigma(D_a)$, then

$$(\sigma_{-a'}\cup\sigma_{-a})\ \#\ (\sigma(L)\#\sigma(L'))\ \#\ (\sigma_{-b'}\cup\sigma_{-b}) = [\sigma_{-a'}\cup(\sigma_{-a}\#\sigma(L))]\ \#\ [(\sigma(L')\#\sigma_{-b'})\cup\sigma_{-b}].$$

The latter is a gluing of all ends, so we can apply the first claim provided that the ends are correctly ordered. Apply Lemma 103 iteratively to correctly reorder the (negative) ends of the second square bracket: this gives rise to the sign in the second claim. Except for this sign, the result of the gluing must be the complex orientation just like for $(\sigma_{-a'}\cup\sigma_{-a})\#(\sigma(L\#L'))\#(\sigma_{-b'}\cup\sigma_{-b})$. Removing the caps that we added in both cases shows $\sigma(L)\#\sigma(L')$, $\sigma(L\#L')$ differ by that sign. □

16.13. **Orientation signs for the TQFT.** Now we define coherent orientations for the moduli spaces $\mathcal{M}(x_a;y_b;S,\beta)$. Pick once and for all a choice of trivialization of $x^*T\overline{M}$ over each Hamiltonian orbit x (see 16.16 to avoid choices).

For $u\in\mathcal{M}(x_a;y_b;S,\beta)$, the linearization $L=D_u$ of $\overline{\partial}$ from 15.7 is an operator of the above type on $E=u^*T\overline{M}\to S$ with asymptotics $(L_1^-,\dots,L_p^-;L_1^+,\dots,L_q^+)$ in the chosen trivializations over the asymptotics x_a, y_b. Define the orientation sign $\epsilon_u\in\{\pm1\}$ arising in 15.10 by (notice the order!)

$$\epsilon_u = \sigma(u) = \sigma(\mathcal{O}_{u^*T\overline{M}\to S}(L_1^-,\dots,L_p^-;L_q^+,\dots,L_1^+)).$$

Note that the definition involves the bundle $u^*T\overline{M}$, not just the asymptotics $L^\pm$.

16.14. **Using orientation signs to prove TQFT maps compose correctly.** By coherence, $\sigma(u\#v) = \pm\sigma(u)\#\sigma(v)$ for any broken Floer solution $u\#v$, where the sign $\pm$ is determined by applying Lemma 104. It is $+$ if all the positive ends of u get glued, or if all the negative ends of v get glued. In Theorem 91, $\psi_S\circ\psi_{S'} = \psi_{S\#_\lambda S'}$ for a large λ, since the unique glued solution $u\#_\lambda v$ close to an isolated broken solution $u\#v$ is counted with sign $\sigma(u\#_\lambda v) = \sigma(u\#v) = \sigma(u)\#\sigma(v)$ by coherence.

16.15. **Using orientation signs to prove TQFT maps are chain maps.** If SH^* is $\mathbb{Z}$-graded (2.6), the differential ∂ on $SC^*(H_1)\otimes\cdots SC^*(H_k)$ is defined by

$$\begin{aligned}\partial(a_1\otimes a_2\otimes a_3\otimes\cdots) &= d(a_1)\otimes a_2\otimes a_3\otimes\cdots+(-1)^{|a_1|}a_1\otimes d(a_2)\otimes a_3\otimes\cdots+\\ &\quad+(-1)^{|a_1|+|a_2|}a_1\otimes a_2\otimes d(a_3)\otimes\cdots+\cdots,\end{aligned}$$

this is the Koszul sign convention with d in degree 1. In general, SH^* is only $\mathbb{Z}/2\mathbb{Z}$-graded, but this ∂ makes sense since only the parity of $|a_1|, |a_2|, \ldots$ matter.

Suppose $u\#v$ is a broken Floer solution, where an isolated Floer trajectory u broke off at the first negative end. The gluing of the linearizations $D_u\#D_v$ is

$$\mathcal{O}(D_u^-,D_u^+)\times\mathcal{O}(D_u^+,L_2^-,\ldots,L_p^-;L_q^+,\ldots,L_1^+)\to\mathcal{O}(D_u^-,L_2^-,\ldots,L_p^-;L_q^+,\ldots,L_1^+),$$

so by Lemma 104, $\sigma(u\#v)=\sigma(u)\#\sigma(v)$. Similarly, for broken Floer solutions $v\#u$ when u breaks off at the first positive end. For all other breakings, first apply Lemma 103 to reorder the ends so that we are in one of the previous two situations, then glue, and finally reorder the ends back to their original position. Since we glued an isolated Floer trajectory, the grading of the orbit at that end changes parity, thus the sign introduced by the second reordering differs from the sign of the first reordering precisely by the Koszul sign for ∂ discussed above.

Thus we proved that $\sigma(u)\#\sigma(v)$ and $\sigma(u\#v)$ differ by the Koszul signs that arise in the computation of $\partial\circ\psi_S$, and similarly for $v\#u$ and $\psi_S\circ\partial$. Now repeat verbatim the gluing description of 16.7, replacing the continuation solution v by the Floer solution v, and $\mathcal{M}\subset\mathcal{M}_1(x_a;y_b;S,\beta)$. The gluing map sends $\partial_s u$ to an inward/outward pointing vector near the boundary for breakings of type $v\#u$, $u\#v$ respectively. Thus we obtain the required signs that prove $\psi_S\circ\partial-\partial\circ\psi_S=0$.

16.16. **The choice of trivializations over the Hamiltonian orbits.** In 16.13 we proved that the moduli spaces $\mathcal{M}(x_a;y_b;S,\beta)$ are canonically oriented *relative to the ends*: we chose trivializations of $x^*T\overline{M}$ over the asymptotics $x=x_a,y_b$. Since we are taking determinants, we only need to choose an orientation of $x^*T\overline{M}$. One can incorporate these choices into the symplectic chain complex:

For a 1-dimensional $\mathbb{R}$-vector space V, define the *orientation space* $\mathcal{O}(V)$ to be the 1-dimensional $\mathbb{K}$-vector space generated by the two possible orientations of V, subject to the relation that the two generators sum to zero. This is functorial: any $\mathbb{R}$-linear map $V\to W$ defines a natural $\mathbb{K}$-linear map $\mathcal{O}(V)\to\mathcal{O}(W)$, and similarly for multi-linear maps. In our setup, let $\mathcal{O}(x)$ be generated by the two orientations of $x^*T\overline{M}$ (modulo the relation that they are opposite generators). Then define

$$SC^*(H)=\bigoplus\left\{\mathcal{O}(x): x\in\mathcal{L}\overline{M},\ \dot{x}(t)=X(x(t))\right\}.$$

One then checks that the orientation signs we constructed for the moduli spaces $\mathcal{M}(x,y)$ and $\mathcal{M}(x_a;y_b;S,\beta)$ depend multiplicatively on the choices of orientation of $T\overline{M}$ over the asymptotics. For example, $u\in\mathcal{M}_0(x,y)$ defines an isomorphism $\epsilon_u:\mathcal{O}(y)\to\mathcal{O}(x)$, which, if we made explicit choices of generators, would be multiplication by the sign ϵ_u of 2.1. Then we define the differential d acting on the summand $\mathcal{O}(y)$ by the linear map $\sum\epsilon_u$, summing over $u\in\mathcal{M}_0(x,y)$.

Floer solutions $u\in\mathcal{M}_0(x_a;y_b;S,\beta)$ yield isomorphisms $\epsilon_u:\otimes_b\mathcal{O}(y_b)\to\otimes_a\mathcal{O}(x_a)$, and the operations ψ_S act on $\otimes_b\mathcal{O}(y_b)$ via $\sum\epsilon_u$, summing over all such u.

For twisted symplectic cohomology, we replace the field $\mathbb{K}$ by the Novikov algebra Λ, and we insert the weights $t^{\alpha[u]}$ in front of the ϵ_u maps.

We recommend Seidel [34, Section 12f] for more information about orientations.

17. Appendix 3: Using non-linear Hamiltonians.

17.1. **An alternative definition of $SH^*(M)$.** Lemma 105 shows that we could define $SH^*(M) = SH^*(Q)$. We call *non-linear Hamiltonians* the Hamiltonians Q of the Lemma, such as $\frac{1}{2}R^2$ (one could easily weaken the conditions on Q).

Lemma 105. *Let $Q : \overline{M} \to \mathbb{R}$ equal a convex function $q(R)$ on the collar with $q'(R) \to \infty$. Let $H^m = Q$ where $q' \leq m$ and extend by slope m elsewhere. Then*

$$SH^*(Q) = \bigcup_m SH^*(H^m) \cong \varinjlim SH^*(H) = SH^*(M).$$

Proof. By the convexity of q, $\partial_R(-Rq'(R) + q(R)) = -q''(R) \leq 0$, so the action A_Q decreases on the collar. Thus no Floer trajectories have $R(x_-) > R(x_+)$, otherwise they would have negative energy. So for $m \leq m'$ we get inclusions of subcomplexes $SC^*(H^m) \subset SC^*(H^{m'}) \subset SC^*(Q)$. Since $SC^*(Q)$ consists of *finite* linear combinations of generators, the claim follows. ☐

17.2. **Contact type J make transversality fail.** Defining $SH^*(M) = SH^*(Q)$ avoids direct limits, but makes transversality or compactness proofs harder. To achieve transversality one needs to be able to perturb J on all of $\overline{M}$ and in all directions. To guarantee compactness, one needs the maximum principle, so one needs the contact type condition on J at infinity, $J\partial_r = \mathcal{R}$. This leaves no freedom to perturb J in the plane span$(\partial_r, \mathcal{R})$: for example for $\overline{M} = \mathbb{C}$, J is uniquely determined at infinity. So one cannot naively achieve transversality and compactness at the same time. Thus, one needs to allow J to be of non-contact type and one needs to reprove compactness. We found the following trick:

Theorem 106. *Given J of contact type at infinity, there exists a perturbation J' of J, no longer of contact type at infinity, such that J'-Floer solutions stay within compact sets determined by their ends.*

In fact, there are reals $d_n = d_n(J) > 0$ such that a generic perturbation J' of J will work, subject only to the condition that $|J' - J| < d_n$ on $n \leq R \leq n+1$ for all $n \in \mathbb{N}$. (Generic is always in the sense of Baire second category)

Proof. By 15.7 a generic perturbation of J by an arbitrarily small amount achieves transversality for Floer solutions lying in a given compact $W \subset \overline{M}$. However, we cannot naively perturb J inductively on larger and larger compacts, since the maximum principle will no longer hold on these compacts. Even after fixing the asymptotics A, new Floer solutions may appear reaching into the regions where we perturbed J. Thus, we need to have control over the following value k_A as we perturb J inductively: given asympototic data A (Hamiltonian orbits), let

$$k_A(J) = \max\{R(u) : u \text{ is a } J\text{-Floer solution with asymptotic data } A\},$$

and let $K_A(J) = \{R \leq k_A(J)\} \subset \overline{M}$. Note that $k_A(J) < \infty$ by the maximum principle, since J is of contact type for large R. For simplicity, assume J is of contact type for $R \geq 1$ (the argument can easily be adapted to the general case).

Key claim. *Given any $\varepsilon_n > 0$, there exit $d_n > 0$ such that for any sequence of almost complex structures J_n, with $J_0 = J$, satisfying*

$$J_n = J \text{ on } R > n \text{ and } |J_{n+1} - J| < d_m \text{ on } m \leq R \leq m+1, \text{ all } m \leq n$$

we can guarantee that $k_A(J_{n+1}) - k_A(J_n) < \varepsilon_n$ ("the k_A don't increase too much").

To prove the Key claim, suppose by contradiction that this fails at step $n = N + 1$.

Let $J' = J_N$. Then there exists a sequence J'_m converging to J', with $J'_m = J$ on $R > N+1$, and there are J'_m-Floer solutions u_m with asymptotics in A, which leave the compact $K_A(J')$ by at least a distance $\varepsilon_N > 0$.

The u_m lie in the compact $R \leq N+1$ by the maximum principle (on $R > N+1$, $J'_m = J$ is of contact type). Thus by Gromov compactness, a subsequence of the (u_m, J_m) converges to a J'-Floer solution u with asymptotics A.

For simplicity, assume u is not broken (the argument can easily be adapted to the broken case, for example by allowing broken solutions in the definition of k_A). By the convergence, u must also leave the compact $K_A(J')$ by at least a distance $\varepsilon_N > 0$. Thus u contradicts the definition of $k_A(J')$. The Key claim follows.

Pick a sequence $\varepsilon_n > 0$ so that $\varepsilon = \sum \varepsilon_n > 0$ is as small as we desire. Then inductively construct J_{n+1} by making a generic small perturbation of J_n on $R \leq n$, ensuring that J_n is regular for J_n-Floer solutions lying in $R \leq n$. If we make these perturbations in a controlled way, we can ensure that the conditions of the Key claim are met for ε_n, and also that the J_n converge to a regular almost complex structure J'. By construction $k_A(J') < k_A(J) + \varepsilon < \infty$, so the Theorem follows. □

Corollary 107. *Transversality and compactness can both be achieved for the moduli spaces of Floer solutions for non-linear Hamiltonians, after a generic (non-contact type) perturbation of a given J of contact type at infinity (Generic as in 106).*

17.3. **TQFT structure using non-linear Hamiltonians.** By Theorem 106, we can define TQFT operations for the non-linear Hamiltonian Q. Because of the weights, this will involve tensor products of $SH^*(c\,Q)$ for some weights c. However, there is a natural way to identify $SH^*(c\,Q) \cong SH^*(Q)$: first use the Liouville flow so that the pull-back of the Hamiltonian $c\,Q$ becomes $\leq Q$, then apply a continuation map. This defines TQFT operations $SH^*(Q)^{\otimes q} \to SH^*(Q)^{\otimes p}$. Apply Lemma 105:

Corollary 108. *The TQFT structure on $SH^*(Q)$ agrees with the TQFT structure on $SH^*(M) = \varinjlim SH^*(H)$ (taken over linear Hamiltonians).*

Remark 109. *Analogous arguments hold in the wrapped case: for a non-linear Hamiltonian Q, $HW^*(L) \cong HW^*(L;Q)$ and the TQFT structures agree.*

REFERENCES

[1] A. Abbondandolo, M. Schwarz, *On the Floer homology of cotangent bundles*, Comm. Pure Appl. Math. 59, 254–316, 2006.

[2] A. Abbondandolo, M. Schwarz, *Floer homology of cotangent bundles and the loop product*, preprint arXiv:0810.1995v4, 2009.

[3] M. Abouzaid, P. Seidel, *An open string analogue of Viterbo functoriality*, preprint arXiv: 0712.3177v2, 2009.

[4] M. Atiyah, *Topological quantum field theories*, Inst. Hautes Études Sci. Publ. Math. No. 68, 175–186, 1988.

[5] M. Betz, R. L. Cohen, *Graph moduli spaces and cohomology operations*, Turkish J. Math. 18, 23–41, 1994.

[6] F. Bourgeois, K. Mohnke, *Coherent orientations in symplectic field theory*, Math. Z. 248, no. 1, 123–146, 2004.

[7] F. Bourgeois, A. Oancea, *An exact sequence for contact- and symplectic homology*, Invent. Math. 175, no. 3, 611–680, 2009.

[8] G. E. Bredon, *Sheaf Theory*, 2nd ed., GTM 170, Springer, 1997.

[9] M. Chas, D. Sullivan, *String topology*, preprint arXiv:math/9911159, 1999.

[10] K. Cieliebak, *Handle attaching in symplectic homology and the chord conjecture*, J. Eur. Math. Soc. (JEMS) 4, 115–142, 2002.

[11] K. Cieliebak, U. Frauenfelder, *A Floer homology for exact contact embeddings*, Pacific J. Math. 239, no. 2, 251–316, 2009.
[12] K. Cieliebak, U. Frauenfelder, A. Oancea, *Rabinowitz Floer homology and symplectic homology*, arXiv:0903.0768v1, 2009.
[13] R. L. Cohen, M. Schwarz, *A Morse theoretic description of string topology*, preprint arXiv:0809.0868v2, 2008.
[14] Y. Eliashberg, A. Givental, H. Hofer, *Introduction to symplectic field theory*, GAFA 2000 (Tel Aviv, 1999). Geom. Funct. Anal. 2000, Special Volume, Part II, 560–673, 2000.
[15] L. C. Evans, *Partial Differential Equations*, AMS, Graduate Studies in Math. v. 19, 1998.
[16] A. Floer, *Morse Theory for Lagrangian intersections*, J. Diff. Geom. 28, 513–547, 1988.
[17] A. Floer, H. Hofer, *Coherent orientations for periodic orbit problems in symplectic geometry*, Math. Z. 212, no. 1, 13–38, 1993.
[18] A. Floer, H. Hofer, D. Salamon, *Transversality in elliptic Morse theory for the symplectic action*, Duke Math. J. 80, 251–292, 1995.
[19] K. Fukaya, *Morse homotopy and its quantization*, Geometric topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2, AMS, Providence, RI, 409–440, 1997.
[20] K. Fukaya, P. Seidel, I. Smith, "The symplectic geometry of cotangent bundles from a categorical viewpoint" in Homological Mirror Symmetry: New Developments and Perspectives, Lecture Notes in Phys. 757, Springer, Berlin, 1–26, 2009.
[21] V. Guillemin, S. Sternberg, *Supersymmetry and Equivariant De Rham Theory*, Springer, 1999.
[22] M. McLean, *Lefschetz fibrations and symplectic homology*, Geom. Topol. 13, no. 4, 1877–1944, 2009.
[23] D. McDuff, D. Salamon, *J-holomorphic curves and quantum cohomology*, University Lecture Series, 6, AMS, Providence, RI, 1994.
[24] D. McDuff, D. Salamon, *J-holomorphic curves and symplectic topology*, AMS, Colloquim Publications, vol. 52, 2004.
[25] K. Mohnke, *Holomorphic disks and the Chord Conjecture*, Ann. of Math. 154, 219–222, 2001.
[26] S. Piunikhin, D. Salamon, M. Schwarz, *Symplectic Floer-Donaldson theory and quantum cohomology*, Contact and symplectic geometry (Cambridge, 1994), 171–200, Publ. Newton Inst. 8, Cambridge Univ. Press, Cambridge, 1996.
[27] A. F. Ritter, *Novikov-symplectic cohomology and exact Lagrangian embeddings*, Geom. Topol. 13, no. 2, 943-978, 2009.
[28] A. F. Ritter, *Deformations of symplectic cohomology and exact Lagrangians in ALE spaces*, to appear in Geom. Funct. Anal. (GAFA), 2010.
[29] D. Salamon, *Lectures on Floer homology. Symplectic geometry and topology (Park City, UT, 1997)*, 143–229, IAS/Park City Math. Ser., 7, AMS, Providence, RI, 1999.
[30] D. Salamon, J. Weber, *Floer homology and the heat flow*, Geom. Funct. Anal. (GAFA) 16 (2006), no. 5, 1050–1138.
[31] D. Salamon, E. Zehnder, *Morse theory for periodic solutions of Hamiltonian systems and the Maslov index*, Comm. Pure Appl. Math. 45, no. 10, 1303–1360, 1992.
[32] M. Schwarz, *Cohomology Operations from S^1-Cobordisms in Floer Homology*, Ph.D., ETH Zürich, 1995. Available at: http://www.mathematik.uni-leipzig.de/∼ schwarz/
[33] P. Seidel, *A biased view of symplectic cohomology*, Current Developments in Mathematics, 2006, 211-253, Int. Press, Somerville, MA, 2008.
[34] P. Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zürich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008.
[35] D. Sullivan, *Differential Forms and the Topology of Manifolds*, Manifolds – Tokyo 1973 (Proc. Internat. Conf., Tokyo, 1973), 37–49, Univ. Tokyo Press, Tokyo, 1975.
[36] C. Viterbo, *Functors and computations in Floer homology with applications. I.*, Geom. Funct. Anal. 9, no. 5, 985–1033, 1999.
[37] C. Viterbo, *Exact Lagrange submanifolds, periodic orbits and the cohomology of free loop spaces.*, J. Diff. Geom. 47, no. 3, 420–468, 1997.

Trinity College, Cambridge, CB2 1TQ, England.
E-mail address: `a.f.ritter@dpmms.cam.ac.uk`